

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

## PATENT COOPERATION TREATY

PCT

## NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Assistant Commissioner for Patents  
 United States Patent and Trademark  
 Office  
 Box PCT  
 Washington, D.C. 20231  
 ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

<b>Date of mailing</b> (day/month/year) 26 July 2000 (26.07.00)	
<b>International application No.</b> PCT/US99/28589	<b>Applicant's or agent's file reference</b> BB1264 PCT
<b>International filing date</b> (day/month/year) 02 December 1999 (02.12.99)	<b>Priority date</b> (day/month/year) 03 December 1998 (03.12.98)
<b>Applicant</b> CAHOON, Edgar, B. et al	

1. The designated Office is hereby notified of its election made:


☒ in the demand filed with the International Preliminary Examining Authority on:  
 26 June 2000 (26.06.00)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was  
☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

<b>The International Bureau of WIPO</b> 34, chemin des Colombettes 1211 Geneva 20, Switzerland	<b>Authorized officer</b>  Kiwa Mpay
Facsimile No.: (41-22) 740.14.35	Telephone No.: (41-22) 338.83.38



INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)



Applicant's or agent's file reference BB1264		<b>FOR FURTHER ACTION</b> See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/US99/28589	International filing date (day/month/year) 02/12/1999	Priority date (day/month/year) 03/12/1998	
International Patent Classification (IPC) or national classification and IPC C12N15/53			
Applicant E.I. DU POND DE NEMOURS AND COMPANY et al			

- This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.
- This REPORT consists of a total of 9 sheets, including this cover sheet.
  - ☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☒ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☒ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☒ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand  26/06/2000	Date of completion of this report  27.03.2001
Name and mailing address of the international preliminary examining authority:   European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465	Authorized officer  Page, M  Telephone No. +49 89 2399 7322  

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/US99/28589

## I. Basis of the report

1. This report has been drawn on the basis of *(substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments (Rules 70.16 and 70.17).)*:

### Description, pages:

1-30 as originally filed

### Claims, No.:

1-19 as originally filed

### Drawings, sheets:

1/3-3/3 as originally filed

### Sequence listing part of the description, pages:

1-19 (SEQ ID NOs. 1-17), as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☒ contained in the international application in written form.
- ☒ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/US99/28589

- ☐ the description, pages:
- ☐ the claims, Nos.:
- ☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

*(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)*

6. Additional observations, if necessary:

## II. Priority

1. ☐ This report has been established as if no priority had been claimed due to the failure to furnish within the prescribed time limit the requested:

- ☐ copy of the earlier application whose priority has been claimed.
- ☐ translation of the earlier application whose priority has been claimed.

2. ☐ This report has been established as if no priority had been claimed due to the fact that the priority claim has been found invalid.

Thus for the purposes of this report, the international filing date indicated above is considered to be the relevant date.

3. Additional observations, if necessary:  
**see separate sheet**

## IV. Lack of unity of invention

1. In response to the invitation to restrict or pay additional fees the applicant has:

- ☐ restricted the claims.
- ☐ paid additional fees.
- ☐ paid additional fees under protest.
- ☒ neither restricted nor paid additional fees.

2. ☐ This Authority found that the requirement of unity of invention is not complied and chose, according to Rule 68.1, not to invite the applicant to restrict or pay additional fees.

3. This Authority considers that the requirement of unity of invention in accordance with Rules 13.1, 13.2 and 13.3 is

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/US99/28589

☐ complied with.

☒ not complied with for the following reasons:  
**see separate sheet**

4. Consequently, the following parts of the international application were the subject of international preliminary examination in establishing this report:

☐ all parts.

☒ the parts relating to claims Nos. 1-19 (all partially).

## V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

### 1. Statement

Novelty (N)	Yes: Claims 2, 9, 12, 14, 15 (all partially)
	No: Claims 1, 3-8, 10, 11, 13, 16-19 (all partially)
Inventive step (IS)	Yes: Claims
	No: Claims 1-19 (all partially)
Industrial applicability (IA)	Yes: Claims 1-19 (all partially)
	No: Claims

2. Citations and explanations  
**see separate sheet**

## VII. Certain defects in the international application

The following defects in the form or contents of the international application have been noted:  
**see separate sheet**

## VIII. Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:  
**see separate sheet**

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/US99/28589

The application concerns the provision of polynucleotide and polypeptide sequences corresponding to plant (Florida bitterbush) delta-6- or sphingolipid-desaturase. One of the principle products of this enzyme, gamma-linoleic acid, is reported to have pharmaceutical properties.

**Re Item II**

**Priority**

The priority document available to the IPEA at the time of examination only discloses 2 polypeptide sequences (SEQ ID NOs. 4 and 5 of the priority document, the former corresponding to SEQ ID NO. 2 of the present application), but no polynucleotide sequences, although reference is made to such in the document. For now, it has been assumed that priority is valid as claimed.

**Re Item IV**

**Lack of Unity of Invention**

The applicant has not responded to the invitation to pay additional fees or restrict the claims. Examination of novelty, inventive step and industrial applicability has therefore been restricted to Invention 1) as described in the Invitation, i.e. **claims 1-19 (partially)**, corresponding to isolated polynucleotide and polypeptide sequences encoding delta-6/sphingolipid desaturase and their selection and uses where the sequences are SEQ ID NOs. 1 and 2.

**Re Item V**

**Reasoned statement under Rule 66.2(a)(ii) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement**

- 1) Reference is made to the following documents:  
D1: WO 96 21022 A (RHONE POULENC AGROCHIMIE) 11 July 1996 (1996-07-11)  
D2: OLGA SAYANOVA ET AL: 'Expression of a borage desaturase cDNA

containing an N-terminal cytochrome b5 domain results in the accumulation of high levels of Delta6-desaturated fatty acids in transgenic tobacco' PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, US, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, vol. 94, 15 April 1997 (1997-04-15), pages 4211-4216, XP002106758 ISSN: 0027-8424 cited in the application

D3: WO 98 45461 A (RHONE-POULENC AGROCHIMIE ; LI ZHONGSEN (US); THOMAS TERRY L (US)) 15 October 1998 (1998-10-15)

2) **Novelty - Art.33(1) and (2) PCT:**

The novelty of **claims 1 and 10 (partially)** cannot be acknowledged in light of D1, D2 and D3, which disclose the polynucleotide and polypeptide sequences of storage delta-6-desaturases. These enzymes are 55.4% identical to SEQ ID NO. 2 over 444 amino acid residues (D1 SEQ ID NO. 4; D3 SEQ ID NO. 2) and 59.8% identical to SEQ ID NO. 2 over 448 amino acid residues (D2 Fig.1), according to the sequence alignment algorithms used by the IPEA.

**Claims 3-8 (partially)** therefore also lack novelty, as isolated polynucleotides, chimeric genes and transformed isolated host cells are also provided in these documents (D1 page 12 lines 18-19; D1 page 12 line 24 to page 14 line 2; D2 page 4213 Functional Analysis of pBdes6 in Transgenic Tobacco; D3 Examples 9 and 10).

**Claims 2 and 12 (partially)** appear to be novel in light of the cited prior art. The particular sequences disclosed therein do not appear to be found in the prior art.

**Claim 9 (partially)** also appears to be novel in light of the cited prior art, as viruses comprising the nucleotides of the claimed invention are not found therein.

The methods of **claims 11, 13 and 16-19 (partially)** lack novelty. D1 discloses the introduction of such polynucleotides into plants and subsequent assays which determine the presence of the polypeptide in comparison to untransformed plants via the identification of the products of the enzyme (D1 page 6 line 16 to page 7 line 25). D2 also discloses such methods (D2 page 4213 Functional Analysis of pBdes6 in



Transgenic Tobacco).

**Claims 14 and 15 (partially)** also lack novelty. Both D1 and D2 disclose methods for obtaining nucleic acid fragments using amplification (D2 page 4212 PCR-Based Cloning) or hybridisation (D1 page 10 lines 6-12; D2 page 4212 Library Screening and pages 4212-4213 PCR-Based Cloning of Membrane-Bound Desaturases).

The novelty of the above claims could be restored if they were limited to the exact sequences as set forth in SEQ ID NOs. 1 and 2.

3) **Inventive Step - Art.33(1) and (3) PCT:**

The following comments on inventive step are confined to subject matter which could be acknowledged as being novel, or for which novelty could potentially be restored as outlined supra.

The closest prior art is document D1, which discloses the polypeptide and polynucleotide sequences of borage delta-6-desaturase, vectors and host cells comprising such sequences, methods for selecting cells comprising such polynucleotides and methods for obtaining nucleic acids by hybridisation (D1 SEQ ID NO. 4, page 10 lines 6-12, page 12 lines 18-19, page 12 line 24 to page 14 line 2).

In light of the prior art, the objective technical problem can be regarded as the provision of alternative plant delta-6-desaturase polynucleotide and polypeptide sequences.

The technical problem is solved by the subject matter of claims 2, 9 and 12, which provide Florida bitterbrush delta-6-desaturase sequences, a virus comprising a delta-6-desaturase sequence of claim 1 and a method for selecting a nucleotide from SEQ ID NO. 1 encoding a desaturase.

**Claims 2, 9 and 12 (partially)** cannot be considered to demonstrate inventive step. SEQ ID NOs. 1 and 2 are merely alternative sequences to those already known in

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/US99/28589

the art and do not demonstrate any inventive features. In the absence of any beneficial effect, merely providing a further sequence for a known enzyme cannot be regarded as demonstrating inventive step.

**Claims 9 and 12** do not contribute to the teaching in the art and are considered to be trivial.

Even if novelty were to be restored to claims 1, 3-8 and 10-19, it is not possible to see where any inventive step might lie here for the above reasons.

**Re Item VII**

**Certain defects in the international application**

- a) Contrary to the requirements of Rule 5.1(a)(ii) PCT, the relevant background art disclosed in the documents D1 and D3 are not mentioned in the description, nor are these documents identified therein.

**Re Item VIII**

**Certain observations on the international application**

- a) **Claim 8** seeks protection for virus cells. Viruses, however, are not cells.
- b) **Claim 9** lacks support from the description (Articles 5 and 6 PCT). Viruses comprising the isolated polynucleotide of claim 1 are not disclosed therein.
- c) **Claims 11, 14 and 15** lack clarity and the scope of the said claims is unclear (Article 6 PCT). All three claims use the term "comprising a nucleotide sequence of at least one of 30 [or 40 in claim 14] contiguous nucleotides". It is unclear whether the claims seek protection for methods employing oligonucleotides of 1-30 [or 40] nucleotides length or something else.
- d) The selection method of **claims 11-13 and 16-19** is unclear. It is not clear what is to

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/US99/28589

be selected and how it is to be selected. The scope of the said claims is thereby rendered obscure (Article 6 PCT).

- e) Embodiment (c) of **claims 11 and 13** lacks clarity. It is not clear what polypeptide is to be measured (Article 6 PCT).
- f) **Claim 14** lacks clarity as it appears that the claim seeks protection for a method of polynucleotide amplification using only one oligonucleotide primer, unless the claim only seeks protection for first-strand synthesis (Article 6 PCT).

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/28589

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/53 C12N9/02 C12N5/10 G01N33/50 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

STRAND, EPO-Internal, WPI Data, PAJ, BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 21022 A (RHONE-POULENC AGROCHIMIE) 11 July 1996 (1996-07-11) page 1-3 page 9, line 28 -page 11, line 14 SEQ ID NOS 4, 5	1,3,5-8, 10-17,19
A	---	2,4
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

21 August 2000

Date of mailing of the international search report

11. 9. 00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Ceder, O

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/28589

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>OLGA SAYANOVA ET AL: "Expression of a borage desaturase cDNA containing an N-terminal cytochrome b5 domain results in the accumulation of high levels of Delta6-desaturated fatty acids in transgenic tobacco"</p> <p>PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, US, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, vol. 94, 15 April 1997 (1997-04-15), pages 4211-4216, XP002106758</p> <p>ISSN: 0027-8424</p> <p>cited in the application</p> <p>the whole document</p> <p>----</p>	1-8, 10, 14, 15
X	<p>WO 98 45461 A (RHONE POULENC AGROCHIMIE ; LI ZHONGSEN (US); THOMAS TERRY L (US))</p> <p>15 October 1998 (1998-10-15)</p> <p>abstract; claims; examples 2-4, 9, 10</p> <p>----</p>	1, 3, 5-8, 10
A	<p>SPERLING ET AL: "A sphingolipid desaturase from higher plants. Identification of a new cytochrome b5 fusion protein"</p> <p>JOURNAL OF BIOLOGICAL CHEMISTRY, US, AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, vol. 273, no. 44, 30 October 1998 (1998-10-30), pages 28590-28596, XP002132572</p> <p>ISSN: 0021-9258</p> <p>cited in the application</p> <p>----</p>	
A	<p>SHEN ET AL.: "Putative cytochrome B5"</p> <p>EMBL SEQUENCE DATABASE, 1 November 1998 (1998-11-01), XP002145279</p> <p>HEIDELBERG DE</p> <p>Ac 082348</p> <p>the whole document</p> <p>----</p>	10
P, X	<p>DELHAIZE ET AL.: "Triticum aestivum S276" (S276) mRNA, complete cds"</p> <p>EMBL SEQUENCE DATABASE, 6 January 1999 (1999-01-06), XP002145280</p> <p>HEIDELBERG DE</p> <p>Ac AF031194</p> <p>the whole document</p> <p>----</p>	1
P, X	<p>-&amp; DELHAIZE ET AL.: "S276"</p> <p>EMBL SEQUENCE DATABASE, 1 May 1999 (1999-05-01), XP002145315</p> <p>HEIDELBERG DE</p> <p>Ac Q9ZTU8</p> <p>the whole document</p> <p>-----</p>	10

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/28589

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9621022 A	11-07-1996	US 5614393 A	25-03-1997
		AU 707061 B	01-07-1999
		AU 4673596 A	24-07-1996
		BR 9510411 A	19-05-1998
		CA 2207906 A	11-07-1996
		CN 1177379 A	25-03-1998
		EP 0801680 A	22-10-1997
		JP 10511848 T	17-11-1998
		US 5789220 A	04-08-1998
WO 9845461 A	15-10-1998	US 5977436 A	02-11-1999
		AU 7107198 A	30-10-1998
		BR 9807969 A	08-03-2000
		EP 0973920 A	26-01-2000
		ZA 9803047 A	05-03-1999

From the  
INTERNATIONAL PRELIMINARY EXAMINING AUTHORITY

by fax and post

RECEIVED

PCT

MAR 22 2001

NOTIFICATION OF TRANSMITTAL OF  
THE INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT  
(PCT Rule 71.1)

PATENT RECORDS  
CENTER

To:

FEULNER, Gregory, J.  
E.I. Du Pont De Nemours and Company  
Legal Patent Records Center  
1007 Market Street  
Wilmington, DE 19898  
ETATS-UNIS D'AMERIQUE

FAX NO. (302) 773-0164

Date of mailing  
(day/month/year) 27.03.2001

Applicant's or agent's file reference  
BB1264

IMPORTANT NOTIFICATION

International application No.  
PCT/US99/28589

International filing date (day/month/year)  
02/12/1999

Priority date (day/month/year)  
03/12/1998

Applicant

E.I. DU POND DE NEMOURS AND COMPANY et al

1. The applicant is hereby notified that this International Preliminary Examining Authority transmits herewith the international preliminary examination report and its annexes, if any, established on the international application.
2. A copy of the report and its annexes, if any, is being transmitted to the International Bureau for communication to all the elected Offices.
3. Where required by any of the elected Offices, the International Bureau will prepare an English translation of the report (but not of any annexes) and will transmit such translation to those Offices.

4. REMINDER

The applicant must enter the national phase before each elected Office by performing certain acts (filing translations and paying national fees) within 30 months from the priority date (or later in some Offices) (Article 39(1)) (see also the reminder sent by the International Bureau with Form PCT/IB/301).

Where a translation of the international application must be furnished to an elected Office, that translation must contain a translation of any annexes to the international preliminary examination report. It is the applicant's responsibility to prepare and furnish such translation directly to each elected Office concerned.

For further details on the applicable time limits and requirements of the elected Offices, see Volume II of the PCT Applicant's Guide.

Name and mailing address of the IPEA/

European Patent Office  
D-80298 Munich

Tel. +49 89 2399 - 0 Tx: 523656 epmu d  
Fax: +49 89 2399 - 4465

Authorized officer

Büchler, S

Tel. +49 89 2399-8090

REY NOTED



03 JE 2001



## PATENT COOPERATION TREATY

## PCT

## INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)



Applicant's or agent's file reference <b>BB1264</b>	<b>FOR FURTHER ACTION</b> See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. <b>PCT/US99/28589</b>	International filing date (day/month/year) <b>02/12/1999</b>	Priority date (day/month/year) <b>03/12/1998</b>
International Patent Classification (IPC) or national classification and IPC <b>C12N15/53</b>		
Applicant <b>E.I. DU POND DE NEMOURS AND COMPANY et al</b>		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.
2. This REPORT consists of a total of 9 sheets, including this cover sheet.
  - ☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☒ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☒ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☒ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand <b>26/06/2000</b>	Date of completion of this report <b>27.03.2001</b>
Name and mailing address of the international preliminary examining authority:  <b>European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465</b>	Authorized officer  <b>Page, M</b>  Tel phone No. +49 89 2399 7322 

# INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/US99/28589

## I. Basis of the report

1. This report has been drawn on the basis of *(substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments (Rules 70.16 and 70.17).):*

Description, pages:

1-30 as originally filed

Claims, No.:

1-19 as originally filed

Drawings, sheets:

1/3-3/3 as originally filed

Sequence listing part of the description, pages:

1-19 (SEQ ID NOS. 1-17), as originally filed

2. With regard to the language, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☒ contained in the international application in written form.
- ☒ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT**International application No. **PCT/US99/28589**

- ☐ the description,      pages:  
☐ the claims,      Nos.:  
☐ the drawings,      sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):  
*(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)*

6. Additional observations, if necessary:

**II. Priority**

1. ☐ This report has been established as if no priority had been claimed due to the failure to furnish within the prescribed time limit the requested:  
☐ copy of the earlier application whose priority has been claimed.  
☐ translation of the earlier application whose priority has been claimed.
2. ☐ This report has been established as if no priority had been claimed due to the fact that the priority claim has been found invalid.
- Thus for the purposes of this report, the international filing date indicated above is considered to be the relevant date.

3. Additional observations, if necessary:  
**see separate sheet**

**IV. Lack of unity of invention**

1. In response to the invitation to restrict or pay additional fees the applicant has:  
☐ restricted the claims.  
☐ paid additional fees.  
☐ paid additional fees under protest.  
☒ neither restricted nor paid additional fees.
2. ☐ This Authority found that the requirement of unity of invention is not complied and chose, according to Rule 68.1, not to invite the applicant to restrict or pay additional fees.
3. This Authority considers that the requirement of unity of invention in accordance with Rules 13.1, 13.2 and 13.3 is

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT**

International application No. PCT/US99/28589

☐ complied with.

☒ not complied with for the following reasons:  
see separate sheet

4. Consequently, the following parts of the international application were the subject of international preliminary examination in establishing this report:

☐ all parts.

☒ the parts relating to claims Nos. 1-19 (all partially).

**V. Reasoned statement under Article 36(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement**

**1. Statement**

Novelty (N)	Yes: Claims 2, 9, 12, 14, 15 (all partially)
	No: Claims 1, 3-8, 10, 11, 13, 16-19 (all partially)
Inventive step (IS)	Yes: Claims
	No: Claims 1-19 (all partially)
Industrial applicability (IA)	Yes: Claims 1-19 (all partially)
	No: Claims

**2. Citations and explanations**  
see separate sheet

**VII. Certain defects in the international application**

The following defects in the form or contents of the international application have been noted:  
see separate sheet

**VIII. Certain observations on the international application**

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:  
see separate sheet

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/US99/28589

The application concerns the provision of polynucleotide and polypeptide sequences corresponding to plant (Florida bitterbush) delta-6- or sphingolipid-desaturase. One of the principle products of this enzyme, gamma-linoleic acid, is reported to have pharmaceutical properties.

**Re Item II****Priority**

The priority document available to the IPEA at the time of examination only discloses 2 polypeptide sequences (SEQ ID NOs. 4 and 5 of the priority document, the former corresponding to SEQ ID NO. 2 of the present application), but no polynucleotide sequences, although reference is made to such in the document. For now, it has been assumed that priority is valid as claimed.

**Re Item IV****Lack of Unity of Invention**

The applicant has not responded to the invitation to pay additional fees or restrict the claims. Examination of novelty, inventive step and industrial applicability has therefore been restricted to Invention 1) as described in the Invitation, i.e. **claims 1-19 (partially)**, corresponding to isolated polynucleotide and polypeptide sequences encoding delta-6/sphingolipid desaturase and their selection and uses where the sequences are SEQ ID NOs. 1 and 2.

**Re Item V**

**Reasoned statement under Rule 66.2(a)(ii) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement**

- 1) Reference is made to the following documents:  
D1: WO 96 21022 A (RHONE-POULENC AGROCHIMIE) 11 July 1996 (1996-07-11)  
D2: OLGA SAYANOVA ET AL: 'Expression of a borage desaturase cDNA

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/US99/28589

containing an N-terminal cytochrome b5 domain results in the accumulation of high levels of Delta6-desaturated fatty acids in transgenic tobacco' PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, US, NATIONAL ACADEMY OF SCIENCE, WASHINGTON, vol. 94, 15 April 1997 (1997-04-15), pages 4211-4216, XP002106758 ISSN: 0027-8424 cited in the application

D3: WO 98 45461 A (RHONE-POULENC AGROCHIMIE ; LI ZHONGSEN (US); THOMAS TERRY L (US)) 15 October 1998 (1998-10-15)

2) **Novelty - Art.33(1) and (2) PCT:**

The novelty of **claims 1 and 10 (partially)** cannot be acknowledged in light of D1, D2 and D3, which disclose the polynucleotide and polypeptide sequences of borage delta-6-desaturases. These enzymes are 55.4% identical to SEQ ID NO. 2 over 444 amino acid residues (D1 SEQ ID NO. 4; D3 SEQ ID NO. 2) and 59.8% identical to SEQ ID NO. 2 over 448 amino acid residues (D2 Fig.1), according to the sequence alignment algorithms used by the IPEA.

**Claims 3-8 (partially)** therefore also lack novelty, as isolated polynucleotides, chimeric genes and transformed isolated host cells are also provided in these documents (D1 page 12 lines 18-19; D1 page 12 line 24 to page 14 line 2; D2 page 4213 Functional Analysis of pBdes6 in Transgenic Tobacco; D3 Examples 9 and 10).

**Claims 2 and 12 (partially)** appear to be novel in light of the cited prior art. The particular sequences disclosed therein do not appear to be found in the prior art.

**Claim 9 (partially)** also appears to be novel in light of the cited prior art, as viruses comprising the nucleotides of the claimed invention are not found therein.

The methods of **claims 11, 13 and 16-19 (partially)** lack novelty. D1 discloses the introduction of such polynucleotides into plants and subsequent assays which determine the presence of the polypeptide in comparison to untransformed plants via the identification of the products of the enzyme (D1 page 6 line 16 to page 7 line 25). D2 also discloses such methods (D2 page 4213 Functional Analysis of pBdes6 in

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/US99/28589

Transgenic Tobacco).

**Claims 14 and 15 (partially)** also lack novelty. Both D1 and D2 disclose methods for obtaining nucleic acid fragments using amplification (D2 page 4212 PCR-Based Cloning) or hybridisation (D1 page 10 lines 6-12; D2 page 4212 Library Screening and pages 4212-4213 PCR-Based Cloning of Membrane-Bound Desaturases).

The novelty of the above claims could be restored if they were limited to the exact sequences as set forth in SEQ ID NOs. 1 and 2.

3) **Inventive Step - Art.33(1) and (3) PCT:**

The following comments on inventive step are confined to subject matter which could be acknowledged as being novel, or for which novelty could potentially be restored as outlined supra.

The closest prior art is document D1, which discloses the polypeptide and polynucleotide sequences of borage delta-6-desaturase, vectors and host cells comprising such sequences, methods for selecting cells comprising such polynucleotides and methods for obtaining nucleic acids by hybridisation (D1 SEQ ID NO. 4, page 10 lines 6-12, page 12 lines 18-19, page 12 line 24 to page 14 line 2).

In light of the prior art, the objective technical problem can be regarded as the provision of alternative plant delta-6-desaturase polynucleotide and polypeptide sequences.

The technical problem is solved by the subject matter of claims 2, 9 and 12, which provide Florida bitterbrush delta-6-desaturase sequences, a virus comprising a delta-6-desaturase sequence of claim 1 and a method for selecting a nucleotide from SEQ ID NO. 1 encoding a desaturase.

**Claims 2, 9 and 12 (partially)** cannot be considered to demonstrate inventive step. SEQ ID NOs. 1 and 2 are merely alternative sequences to those already known in

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/US99/28589

the art and do not demonstrate any inventive features. In the absence of any beneficial effect, merely providing a further sequence for a known enzyme cannot be regarded as demonstrating inventive step.

**Claims 9 and 12** do not contribute to the teaching in the art and are considered to be trivial.

Even if novelty were to be restored to claims 1, 3-8 and 10-19, it is not possible to see where any inventive step might lie here for the above reasons.

**Re Item VII**

**Certain defects in the international application**

- a) Contrary to the requirements of Rule 5.1(a)(ii) PCT, the relevant background art disclosed in the documents D1 and D3 are not mentioned in the description, nor are these documents identified therein.

**Re Item VIII**

**Certain observations on the international application**

- a) **Claim 8** seeks protection for virus cells. Viruses, however, are not cells.
- b) **Claim 9** lacks support from the description (Articles 5 and 6 PCT). Viruses comprising the isolated polynucleotide of claim 1 are not disclosed therein.
- c) **Claims 11, 14 and 15** lack clarity and the scope of the said claims is unclear (Article 6 PCT). All three claims use the term "comprising a nucleotide sequence of at least one of 30 [or 40 in claim 14] contiguous nucleotides". It is unclear whether the claims seek protection for methods employing oligonucleotides of 1-30 [or 40] nucleotides length or something else.
- d) The selection method of **claims 11-13 and 16-19** is unclear. It is not clear what is to



**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/US99/28589

be selected and how it is to be selected. The scope of the said claims is thereby rendered obscure (Article 6 PCT).

- e) Embodiment (c) of claims 11 and 13 lacks clarity. It is not clear what polypeptide is to be measured (Article 6 PCT).
- f) Claim 14 lacks clarity as it appears that the claim seeks protection for a method of polynucleotide amplification using only one oligonucleotide primer, unless the claim only seeks protection for first-strand synthesis (Article 6 PCT).



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : C12N 15/53, 9/02, 5/10, G01N 33/50, C12Q 1/68		A2	(11) International Publication Number: WO 00/32790
			(43) International Publication Date: 8 June 2000 (08.06.00)
(21) International Application Number: PCT/US99/28589		(81) Designated States: AE, AL, AU, BA, BB, BG, BR, CA, CN, CR, CU, CZ, DM, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 2 December 1999 (02.12.99)			
(30) Priority Data: 60/110,784                      3 December 1998 (03.12.98)                      US			
(71) Applicant (for all designated States except US): E.I. DU PONT DE NEMOURS AND COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): CAHOON, Edgar, B. [US/US]; 2331 West 18th Street, Wilmington, DE 19806 (US). CAHOON, Rebecca, E. [US/US]; 2331 West 18th Street, Wilmington, DE 19806 (US). HITZ, William, D. [US/US]; 404 Hillside Road, Wilmington, DE 19807 (US). KINNEY, Anthony, J. [GB/US]; 609 Lore Avenue, Wilmington, DE 19809 (US).			
(74) Agent: FEULNER, Gregory, J.; E.I. Du Pont De Nemours and Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).		Published Without international search report and to be republished upon receipt of that report.	

**(54) Title:** MEMBRANE-BOUND DESATURASES

**(57) Abstract**

This invention relates to an isolated nucleic acid fragment encoding a delta-6 desaturase or sphingolipid desaturase. The invention also relates to the construction of a chimeric gene encoding all or a portion of the delta-6 desaturase or sphingolipid desaturase, in sense or antisense orientation, wherein expression of the chimeric gene results in production of altered levels of the delta-6 desaturase or sphingolipid desaturase in a transformed host cell.

[illegible]

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## TITLE

### MEMBRANE-BOUND DESATURASES

This application claims the benefit of U.S. Provisional Application No. 60/110,784 filed December 3, 1998.

## FIELD OF THE INVENTION

This invention is in the field of plant molecular biology. More specifically, this invention pertains to nucleic acid fragments encoding delta-6 desaturase or sphingolipid desaturase in plants and seeds.

## BACKGROUND OF THE INVENTION

Polyunsaturated fatty acids are of major importance in animal health as they have roles in the maintenance of membrane structure and function, in the regulation of cholesterol synthesis and transport, in the prevention of water loss from the skin, and as precursors of eicosanoids, including prostaglandins and leucotrienes. In animals, members of this class of fatty acids are synthesized from the essential fatty acid linoleic acid (C18:2  $\Delta^{9,12}$ ), the first step being the desaturation to gamma-linolenic acid (GLA; C18:3  $\Delta^{6,9,12}$ ) catalyzed by delta-6 desaturase. Clinical trials have shown that dietary supplementation of GLA may be effective in treating a number of ailments (e.g., atopic eczema, mastalgia, diabetic neuropathy, viral infections, and some types of cancer) (Jiang et al. (1998) *Crit. Rev. Oncol. Hematol.* 27:179-209; Kruger, and Horrobin (1997) *Prog. Lipid Res.* 36:131-51). Oils containing GLA are therefore widely used as a general health supplement and have been registered for pharmaceutical use.

In the plant kingdom, GLA is an uncommon fatty acid. Major commercial sources of GLA are evening primroses (*Oenothera* spp.), in which GLA accounts for about 8 to 10% of the seed oil, and borage (starflower; *Borago officinalis*) wherein seeds contain some 20 to 25% GLA. These plants, however, suffer from poor agronomic performance and low yield. There is therefore considerable interest in increasing the GLA content of existing crops and in producing GLA in a conventional oil crop. Expression of a cDNA encoding the delta-6 fatty acid desaturase from developing seeds of borage in transgenic tobacco plants resulted in accumulation of GLA and octadecatetraenoic acid (C18:4  $\Delta^{6,9,12,15}$ ) to levels of 13.2% and 9.6% of the total fatty acids, respectively. The borage delta-6 fatty acid desaturase differs from other previously characterized higher plant desaturase enzymes by the presence of an N-terminal domain related to cytochrome b5 (Sayanova et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:4211-4216). This desaturase does not appear to have an N-terminal cleavable endoplasmic reticulum-targeting signal, but the hydrophobic regions present in the protein would be sufficient to allow it to associate with the endomembrane system. A tripartite motif containing eight conserved histidines has been identified in almost all membrane desaturases (HX<sub>(3-4)</sub>HX<sub>(7-41)</sub>HX<sub>(2-3)</sub>HHX<sub>(61-189)</sub>HX<sub>(2-3)</sub>HH). In *Anabaena* and borage delta-6 desaturase a glutamine residue replaces the first histidine of the third element.

Membrane and reserve lipids of plants contain fatty acids with different degrees of unsaturation which are controlled by different desaturase enzymes. A cDNA isolated from ripening sunflower embryos encodes a protein with the conserved three histidine domains characteristic of membrane-bound desaturases. This cDNA also encodes a fusion protein composed of an N-terminal cytochrome b5 and a domain similar to membrane-bound acyl lipid desaturases (Sperling et al. (1995) *Eur. J. Biochem.* 232:798-805). Expression of homologous cDNAs from *Brassica napus* and *Arabidopsis thaliana* in *Saccharomyces cerevisiae* results in significant proportions of new  $\Delta^{8,9}$ -cis/trans-phytosphingenines that accompany the residual C<sub>18</sub>-phytosphinganine predominating in wild-type yeast cells.

These genes encode new members of the cytochrome b5 superfamily which function as a stereounselective sphingolipid desaturase and show trans-activity (Sperling et al. (1998) *J. Biol. Chem.* 273: 28590-28596).

The enzymes encoded by the picramnia cDNAs included in this application possibly catalyze the formation of a triple bond since picramnia seeds accumulate large amounts of tariric acid (18:1 delta-6-acetylenic), an unusual fatty acid that has a triple bond (or acetylenic bond) at the delta-6 carbon. Tariric acid has many of the same industrial uses ascribed to petroselinic acid, thus these cDNAs should be useful in the production of novel fatty acids in the seed oils of transgenic plants. Although with similarities to the delta-6 desaturase, the enzymes encoded by the corn, soybean, and wheat sequences described herein are sphingolipid desaturases since these plants do not produce delta 6 double bonds like picramnia and borage.

### SUMMARY OF THE INVENTION

The present invention relates to isolated polynucleotides comprising a nucleotide sequence encoding a first polypeptide of at least 60 amino acids that has at least 55% identity based on the Clustal method of alignment when compared to a Florida bitterbush delta-6 desaturase polypeptide of SEQ ID NO:2, and an isolated polynucleotide comprising a nucleotide sequence encoding a first polypeptide of at least 114 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of a corn sphingolipid desaturase polypeptide of SEQ ID NO:4, a soybean sphingolipid desaturase polypeptide of SEQ ID NOs:6 and 8, a wheat sphingolipid desaturase polypeptide of SEQ ID NO:10 and SEQ ID NO:17. The present invention also relates to an isolated polynucleotide comprising the complement of the nucleotide sequences described above.

It is preferred that the isolated polynucleotides of the claimed invention consists of a nucleic acid sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, and 16, that codes for the polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8, and 17. The present invention also relates to an isolated polynucleotide comprising a nucleotide sequences of at least one of 60 (preferably at least one of 40, most preferably at

least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, and 16, and the complement of such nucleotide sequences.

The present invention relates to a chimeric gene comprising an isolated polynucleotide of the present invention operably linked to suitable regulatory sequences.

The present invention relates to an isolated host cell comprising a chimeric gene of the present invention or an isolated polynucleotide of the present invention. The host cell may be eukaryotic, such as a yeast or a plant cell, or prokaryotic, such as a bacterial cell. The present invention also relates to a virus, preferably a baculovirus, comprising an isolated polynucleotide of the present invention or a chimeric gene of the present invention.

The present invention relates to a process for producing an isolated host cell comprising a chimeric gene of the present invention or an isolated polynucleotide of the present invention, the process comprising either transforming or transfecting an isolated compatible host cell with a chimeric gene or isolated polynucleotide of the present invention.

The present invention relates to a delta-6 desaturase polypeptide of at least 60 amino acids comprising at least 55% homology based on the Clustal method of alignment compared to the polypeptide of SEQ ID NO:2, or a sphingolipid desaturase polypeptide of at least 114 amino acids comprising at least 80% homology based on the Clustal method of alignment compared to a polypeptide selected from the group consisting of SEQ ID NOs:4, 6, 8, and 17.

The present invention relates to a method of selecting an isolated polynucleotide that affects the level of expression of a delta-6 desaturase or sphingolipid desaturase polypeptide in a host cell, preferably a plant cell, the method comprising the steps of:

(a) constructing an isolated polynucleotide of the present invention or an isolated chimeric gene of the present invention;

(b) introducing the isolated polynucleotide or the isolated chimeric gene into a host cell;

(c) measuring the level a delta-6 desaturase or sphingolipid desaturase polypeptide in the plant cell containing the isolated polynucleotide; and

(d) comparing the level of a delta-6 desaturase or sphingolipid desaturase polypeptide in the host cell containing the isolated polynucleotide with the level of a delta-6 desaturase or sphingolipid desaturase polypeptide in a host cell that does not contain the isolated polynucleotide.

The present invention relates to a method of obtaining a nucleic acid fragment encoding a substantial portion of a delta-6 desaturase or sphingolipid desaturase polypeptide gene, preferably a plant delta-6 desaturase or sphingolipid desaturase polypeptide gene, comprising the steps of: synthesizing an oligonucleotide primer comprising a nucleotide sequence of at least one of 40 (preferably at least one of 30) contiguous nucleotides derived

from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, and 16, and the complement of such nucleotide sequences; and amplifying a nucleic acid fragment (preferably a cDNA inserted in a cloning vector) using the oligonucleotide primer. The amplified nucleic acid fragment preferably will encode a portion of a delta-6 desaturase or sphingolipid desaturase amino acid sequence.

The present invention also relates to a method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a delta-6 desaturase or sphingolipid desaturase polypeptide comprising the steps of: probing a cDNA or genomic library with an isolated polynucleotide of the present invention; identifying a DNA clone that hybridizes with an isolated polynucleotide of the present invention; isolating the identified DNA clone; and sequencing the cDNA or genomic fragment that comprises the isolated DNA clone.

The present invention also relates to a method for positive selection of a transformed cell comprising:

(a) transforming a host cell with the chimeric gene of the present invention; and  
(b) growing the transformed host cell under conditions suitable for the expression of the chimeric gene allowing expression of the polynucleotide in an amount to alter the concentration of fatty acids with delta-6 double bonds in the host cell to provide a positive selection means.

The present invention also relates to the method of the present invention wherein the host cell is selected from the group consisting of plant cells and procaryotes.

The present invention also relates to the method of the present invention wherein levels of tariric acid are altered.

The present invention also relates to a method for positive selection of a transformed cell comprising:

(a) transforming a plant cell with the chimeric gene of the present invention; and  
(b) growing a plant from the transformed plant cell of step (a) allowing expression of the polynucleotide in an amount to alter the concentration of fatty acids with delta-6 double bonds in the seeds of the plant to provide a positive selection means.

#### BRIEF DESCRIPTION OF THE DRAWINGS AND SEQUENCE DESCRIPTIONS

The invention can be more fully understood from the following detailed description and the accompanying drawings and Sequence Listing which form a part of this application.

Figure 1 shows a comparison of the amino acid sequences of the Florida bitterbush delta-6 desaturase from clone pps.pk0011.d5: fis (SEQ ID NO:2, denoted [SIN 2]), the soybean sphingolipid desaturase from clone ssl.pk0017.b4: fis (SEQ ID NO:8, denoted [SIN 8], the borage delta-6 desaturase having NCBI General Identification No. 2062403 (SEQ ID NO:11), corn sphingolipid desaturase from clone cde1c.pk001.o8: fis (SEQ ID

NO:4, denoted [SIN 4]), wheat sphingolipid desaturase from clone wre1.pk0004.c7: fis (SEQ ID NO:10, denoted [SIN 10],) wheat delta-6 desaturase-like protein having NCBI General Identifier No. 4104056 (SEQ ID NO:12), soybean sphingolipid desaturase from the contig assembled of clones sfl1.pk0012.c5 and sfl1.pk0031.d11 (SEQ ID NO:6, denoted [SIN 6]), and sunflower having NCBI General Identifier No. 1040729 (SEQ ID NO:13).

Table 1 lists the polypeptides that are described herein, the designation of the cDNA clones that comprise the nucleic acid fragments encoding polypeptides representing all or a substantial portion of these polypeptides, and the corresponding identifier (SEQ ID NO:) as used in the attached Sequence Listing. The sequence descriptions and Sequence Listing attached hereto comply with the rules governing nucleotide and/or amino acid sequence disclosures in patent applications as set forth in 37 C.F.R. §1.821-1.825.

**TABLE 1**  
Delta-6 Desaturase or Sphingolipid Desaturase

Delta-6 Fatty Acid Desaturase	Clone Designation	SEQ ID NO:	
		(Nucleotide)	(Amino Acid)
Bitterbush-Fl. [ <i>Picramnia pentandra</i> ]	pps.pk0011.d5: fis	1	2
Sphingolipid Desaturase	Clone Designation	SEQ ID NO:	
		(Nucleotide)	(Amino Acid)
Corn [ <i>Zea mays</i> ]	cde1c.pk001.o8: fis	3	4
Soybean [ <i>Glycine max</i> ]	Contig of: sfl1.pk0012.c5 sfl1.pk0031.d11	5	6
Soybean [ <i>Glycine max</i> ]	ssl.pk0017.b4: fis	7	8
Wheat [ <i>Triticum aestivum</i> ]	wre1.pk0004.c7: fis	9	10
Wheat [ <i>Triticum aestivum</i> ]	wre1.pk0004.c7	16	17

The Sequence Listing contains the one letter code for nucleotide sequence characters and the three letter codes for amino acids as defined in conformity with the IUPAC-IUBMB standards described in *Nucleic Acids Res.* 13:3021-3030 (1985) and in the *Biochemical J.* 219 (No. 2):345-373 (1984) which are herein incorporated by reference. The symbols and format used for nucleotide and amino acid sequence data comply with the rules set forth in 37 C.F.R. §1.822.

#### DETAILED DESCRIPTION OF THE INVENTION

In the context of this disclosure, a number of terms shall be utilized. As used herein, a "polynucleotide" is a nucleotide sequence such as a nucleic acid fragment. A polynucleotide may be a polymer of RNA or DNA that is single- or double-stranded, that optionally contains synthetic, non-natural or altered nucleotide bases. A polynucleotide in the form of



a polymer of DNA may be comprised of one or more segments of cDNA, genomic DNA, synthetic DNA, or mixtures thereof. An isolated polynucleotide of the present invention may include at least one of 60 contiguous nucleotides, preferably at least one of 40 contiguous nucleotides, most preferably one of at least 30 contiguous nucleotides, of the nucleic acid sequence of the SEQ ID NOs:1, 3, 5, 7, 9, 16, or the complement of such sequences.

As used herein, "contig" refers to a nucleotide sequence that is assembled from two or more constituent nucleotide sequences that share common or overlapping regions of sequence homology. For example, the nucleotide sequences of two or more nucleic acid fragments can be compared and aligned in order to identify common or overlapping sequences. Where common or overlapping sequences exist between two or more nucleic acid fragments, the sequences (and thus their corresponding nucleic acid fragments) can be assembled into a single contiguous nucleotide sequence.

As used herein, "substantially similar" refers to nucleic acid fragments wherein changes in one or more nucleotide bases results in substitution of one or more amino acids, but do not affect the functional properties of the polypeptide encoded by the nucleotide sequence. "Substantially similar" also refers to nucleic acid fragments wherein changes in one or more nucleotide bases does not affect the ability of the nucleic acid fragment to mediate alteration of gene expression by gene silencing through for example antisense or co-suppression technology. "Substantially similar" also refers to modifications of the nucleic acid fragments of the instant invention such as deletion or insertion of one or more nucleotides that do not substantially affect the functional properties of the resulting transcript vis-à-vis the ability to mediate gene silencing or alteration of the functional properties of the resulting protein molecule. It is therefore understood that the invention encompasses more than the specific exemplary nucleotide or amino acid sequences and includes functional equivalents thereof.

Substantially similar nucleic acid fragments may be selected by screening nucleic acid fragments representing subfragments or modifications of the nucleic acid fragments of the instant invention, wherein one or more nucleotides are substituted, deleted and/or inserted, for their ability to affect the level of the polypeptide encoded by the unmodified nucleic acid fragment in a plant or plant cell. For example, a substantially similar nucleic acid fragment representing at least one of 30 contiguous nucleotides derived from the instant nucleic acid fragment can be constructed and introduced into a plant or plant cell. The level of the polypeptide encoded by the unmodified nucleic acid fragment present in a plant or plant cell exposed to the substantially similar nucleic fragment can then be compared to the level of the polypeptide in a plant or plant cell that is not exposed to the substantially similar nucleic acid fragment.

For example, it is well known in the art that antisense suppression and co-suppression of gene expression may be accomplished using nucleic acid fragments representing less than the entire coding region of a gene, and by nucleic acid fragments that do not share 100% sequence identity with the gene to be suppressed. Moreover, alterations in a nucleic acid fragment which result in the production of a chemically equivalent amino acid at a given site, but do not effect the functional properties of the encoded polypeptide, are well known in the art. Thus, a codon for the amino acid alanine, a hydrophobic amino acid, may be substituted by a codon encoding another less hydrophobic residue, such as glycine, or a more hydrophobic residue, such as valine, leucine, or isoleucine. Similarly, changes which result in substitution of one negatively charged residue for another, such as aspartic acid for glutamic acid, or one positively charged residue for another, such as lysine for arginine, can also be expected to produce a functionally equivalent product. Nucleotide changes which result in alteration of the N-terminal and C-terminal portions of the polypeptide molecule would also not be expected to alter the activity of the polypeptide. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity of the encoded products. Consequently, an isolated polynucleotide comprising a nucleotide sequence of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, and 16, and the complement of such nucleotide sequences may be used in methods of selecting an isolated polynucleotide that affects the expression of a polypeptide such as a delta-6 fatty acid desaturase and/or a sphingolipid desaturase in a host cell. A method of selecting an isolated polynucleotide that affects the level of expression of a polypeptide in a host cell (eukaryotic, such as plant or yeast, prokaryotic such as bacterial, or viral) may comprise the steps of: constructing an isolated polynucleotide of the present invention or an isolated chimeric gene of the present invention; introducing the isolated polynucleotide or the isolated chimeric gene into a host cell; measuring the level a polypeptide in the host cell containing the isolated polynucleotide; and comparing the level of a polypeptide in the host cell containing the isolated polynucleotide with the level of a polypeptide in a host cell that does not contain the isolated polynucleotide.

Moreover, substantially similar nucleic acid fragments may also be characterized by their ability to hybridize. Estimates of such homology are provided by either DNA-DNA or DNA-RNA hybridization under conditions of stringency as is well understood by those skilled in the art (Hames and Higgins, Eds. (1985) *Nucleic Acid Hybridisation*, IRL Press, Oxford, U.K.). Stringency conditions can be adjusted to screen for moderately similar fragments, such as homologous sequences from distantly related organisms, to highly similar fragments, such as genes that duplicate functional enzymes from closely related organisms. Post-hybridization washes determine stringency conditions. One set of preferred conditions

uses a series of washes starting with 6X SSC, 0.5% SDS at room temperature for 15 min, then repeated with 2X SSC, 0.5% SDS at 45°C for 30 min, and then repeated twice with 0.2X SSC, 0.5% SDS at 50°C for 30 min. A more preferred set of stringent conditions uses higher temperatures in which the washes are identical to those above except for the temperature of the final two 30 min washes in 0.2X SSC, 0.5% SDS was increased to 60°C. Another preferred set of highly stringent conditions uses two final washes in 0.1X SSC, 0.1% SDS at 65°C.

Substantially similar nucleic acid fragments of the instant invention may also be characterized by the percent identity of the amino acid sequences that they encode to the amino acid sequences disclosed herein, as determined by algorithms commonly employed by those skilled in this art. Suitable nucleic acid fragments (isolated polynucleotides of the present invention) encode polypeptides that are at least about 70% identical, preferably at least about 80% identical to the amino acid sequences reported herein. Preferred nucleic acid fragments encode amino acid sequences that are at least about 85% identical to the amino acid sequences reported herein. More preferred nucleic acid fragments encode amino acid sequences that are at least about 90% identical to the amino acid sequences reported herein. Most preferred are nucleic acid fragments that encode amino acid sequences that are at least about 95% identical to the amino acid sequences reported herein. Suitable nucleic acid fragments not only have the above homologies but typically encode a polypeptide having at least about 50 amino acids, preferably at least about 100 amino acids, more preferably at least about 150 amino acids, still more preferably at least about 200 amino acids, and most preferably at least about 250 amino acids. Sequence alignments and percent identity calculations were performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5.

A "substantial portion" of an amino acid or nucleotide sequence comprises an amino acid or a nucleotide sequence that is sufficient to afford putative identification of the protein or gene that the amino acid or nucleotide sequence comprises. Amino acid and nucleotide sequences can be evaluated either manually by one skilled in the art, or by using computer-based sequence comparison and identification tools that employ algorithms such as BLAST (Basic Local Alignment Search Tool; Altschul et al. (1993) *J. Mol. Biol.* 215:403-410; see also [www.ncbi.nlm.nih.gov/BLAST/](http://www.ncbi.nlm.nih.gov/BLAST/)). In general, a sequence of ten or more contiguous amino acids or thirty or more contiguous nucleotides is necessary in order to putatively identify a polypeptide or nucleic acid sequence as homologous to a known protein or gene. Moreover, with respect to nucleotide sequences, gene-specific oligonucleotide probes

comprising 30 or more contiguous nucleotides may be used in sequence-dependent methods of gene identification (e.g., Southern hybridization) and isolation (e.g., *in situ* hybridization of bacterial colonies or bacteriophage plaques). In addition, short oligonucleotides of 12 or more nucleotides may be used as amplification primers in PCR in order to obtain a particular nucleic acid fragment comprising the primers. Accordingly, a “substantial portion” of a nucleotide sequence comprises a nucleotide sequence that will afford specific identification and/or isolation of a nucleic acid fragment comprising the sequence. The instant specification teaches amino acid and nucleotide sequences encoding polypeptides that comprise one or more particular plant proteins. The skilled artisan, having the benefit of the sequences as reported herein, may now use all or a substantial portion of the disclosed sequences for purposes known to those skilled in this art. Accordingly, the instant invention comprises the complete sequences as reported in the accompanying Sequence Listing, as well as substantial portions of those sequences as defined above.

“Codon degeneracy” refers to divergence in the genetic code permitting variation of the nucleotide sequence without effecting the amino acid sequence of an encoded polypeptide. Accordingly, the instant invention relates to any nucleic acid fragment comprising a nucleotide sequence that encodes all or a substantial portion of the amino acid sequences set forth herein. The skilled artisan is well aware of the “codon-bias” exhibited by a specific host cell in usage of nucleotide codons to specify a given amino acid. Therefore, when synthesizing a nucleic acid fragment for improved expression in a host cell, it is desirable to design the nucleic acid fragment such that its frequency of codon usage approaches the frequency of preferred codon usage of the host cell.

“Synthetic nucleic acid fragments” can be assembled from oligonucleotide building blocks that are chemically synthesized using procedures known to those skilled in the art. These building blocks are ligated and annealed to form larger nucleic acid fragments which may then be enzymatically assembled to construct the entire desired nucleic acid fragment. “Chemically synthesized”, as related to nucleic acid fragment, means that the component nucleotides were assembled *in vitro*. Manual chemical synthesis of nucleic acid fragments may be accomplished using well established procedures, or automated chemical synthesis can be performed using one of a number of commercially available machines. Accordingly, the nucleic acid fragments can be tailored for optimal gene expression based on optimization of nucleotide sequence to reflect the codon bias of the host cell. The skilled artisan appreciates the likelihood of successful gene expression if codon usage is biased towards those codons favored by the host. Determination of preferred codons can be based on a survey of genes derived from the host cell where sequence information is available.

“Gene” refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. “Native gene” refers to a gene as found in nature with its

own regulatory sequences. "Chimeric gene" refers any gene that is not a native gene, comprising regulatory and coding sequences that are not found together in nature.

Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived  
5 from the same source, but arranged in a manner different than that found in nature.

"Endogenous gene" refers to a native gene in its natural location in the genome of an organism. A "foreign" gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A "transgene" is a gene  
10 that has been introduced into the genome by a transformation procedure.

"Coding sequence" refers to a nucleotide sequence that codes for a specific amino acid sequence. "Regulatory sequences" refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the  
15 associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

"Promoter" refers to a nucleotide sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. The promoter sequence consists of proximal and more distal upstream  
20 elements, the latter elements often referred to as enhancers. Accordingly, an "enhancer" is a nucleotide sequence which can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise  
25 synthetic nucleotide segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions.

Promoters which cause a nucleic acid fragment to be expressed in most cell types at most times are commonly referred to as "constitutive promoters". New promoters of various  
30 types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamuro and Goldberg (1989) *Biochemistry of Plants* 15:1-82. It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, nucleic acid fragments of different lengths may have identical promoter activity.

35 The "translation leader sequence" refers to a nucleotide sequence located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect processing of the primary transcript to mRNA,

mRNA stability or translation efficiency. Examples of translation leader sequences have been described (Turner and Foster (1995) *Mol. Biotechnol.* 3:225-236).

The "3' non-coding sequences" refer to nucleotide sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. The use of different 3' non-coding sequences is exemplified by Ingelbrecht et al. (1989) *Plant Cell* 1:671-680.

"RNA transcript" refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from posttranscriptional processing of the primary transcript and is referred to as the mature RNA. "Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into polypeptide by the cell. "cDNA" refers to a double-stranded DNA that is complementary to and derived from mRNA. "Sense" RNA refers to an RNA transcript that includes the mRNA and so can be translated into a polypeptide by the cell. "Antisense RNA" refers to an RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene (see U.S. Patent No. 5,107,065, incorporated herein by reference). The complementarity of an antisense RNA may be with any part of the specific nucleotide sequence, i.e., at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. "Functional RNA" refers to sense RNA, antisense RNA, ribozyme RNA, or other RNA that may not be translated but yet has an effect on cellular processes.

The term "operably linked" refers to the association of two or more nucleic acid fragments on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

The term "expression", as used herein, refers to the transcription and stable accumulation of sense (mRNA) or antisense RNA derived from the nucleic acid fragment of the invention. Expression may also refer to translation of mRNA into a polypeptide. "Antisense inhibition" refers to the production of antisense RNA transcripts capable of suppressing the expression of the target protein. "Overexpression" refers to the production of a gene product in transgenic organisms that exceeds levels of production in normal or non-transformed organisms. "Co-suppression" refers to the production of sense RNA transcripts capable of suppressing the expression of identical or substantially similar foreign or endogenous genes (U.S. Patent No. 5,231,020, incorporated herein by reference).

“Altered levels” refers to the production of gene product(s) in transgenic organisms in amounts or proportions that differ from that of normal or non-transformed organisms.

“Mature” protein refers to a post-translationally processed polypeptide; i.e., one from which any pre- or propeptides present in the primary translation product have been removed.

5 “Precursor” protein refers to the primary product of translation of mRNA; i.e., with pre- and propeptides still present. Pre- and propeptides may be but are not limited to intracellular localization signals.

A “chloroplast transit peptide” is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the chloroplast or other plastid types present in the cell in which the protein is made. “Chloroplast transit sequence” refers to a nucleotide sequence that encodes a chloroplast transit peptide. A “signal peptide” is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the secretory system (Chrispeels (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53). If the protein is to be directed to a vacuole, a vacuolar targeting signal (*supra*) can further be added, or if to the endoplasmic reticulum, an endoplasmic reticulum retention signal (*supra*) may be added. If the protein is to be directed to the nucleus, any signal peptide present should be removed and instead a nuclear localization signal included (Raikhel (1992) *Plant Phys.* 100:1627-1632).

20 “Transformation” refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as “transgenic” organisms. Examples of methods of plant transformation include *Agrobacterium*-mediated transformation (De Blaere et al. (1987) *Meth. Enzymol.* 143:277) and particle-accelerated or “gene gun” transformation technology (Klein et al. (1987) *Nature (London)* 327:70-73; U.S. Patent No. 4,945,050, incorporated herein by reference).

Standard recombinant DNA and molecular cloning techniques used herein are well known in the art and are described more fully in Sambrook et al. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, 1989 (hereinafter “Maniatis”).

30 Nucleic acid fragments encoding at least a portion of several delta-6 desaturase or sphingolipid desaturase have been isolated and identified by comparison of random plant cDNA sequences to public databases containing nucleotide and protein sequences using the BLAST algorithms well known to those skilled in the art. The nucleic acid fragments of the instant invention may be used to isolate cDNAs and genes encoding homologous proteins from the same or other plant species. Isolation of homologous genes using sequence-dependent protocols is well known in the art. Examples of sequence-dependent protocols include, but are not limited to, methods of nucleic acid hybridization, and methods of DNA

and RNA amplification as exemplified by various uses of nucleic acid amplification technologies (e.g., polymerase chain reaction, ligase chain reaction).

For example, genes encoding other delta-6 desaturase or sphingolipid desaturases, either as cDNAs or genomic DNAs, could be isolated directly by using all or a portion of the instant nucleic acid fragments as DNA hybridization probes to screen libraries from any desired plant employing methodology well known to those skilled in the art. Specific oligonucleotide probes based upon the instant nucleic acid sequences can be designed and synthesized by methods known in the art (Maniatis). Moreover, the entire sequences can be used directly to synthesize DNA probes by methods known to the skilled artisan such as random primer DNA labeling, nick translation, or end-labeling techniques, or RNA probes using available *in vitro* transcription systems. In addition, specific primers can be designed and used to amplify a part or all of the instant sequences. The resulting amplification products can be labeled directly during amplification reactions or labeled after amplification reactions, and used as probes to isolate full length cDNA or genomic fragments under conditions of appropriate stringency.

In addition, two short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols to amplify longer nucleic acid fragments encoding homologous genes from DNA or RNA. The polymerase chain reaction may also be performed on a library of cloned nucleic acid fragments wherein the sequence of one primer is derived from the instant nucleic acid fragments, and the sequence of the other primer takes advantage of the presence of the polyadenylic acid tracts to the 3' end of the mRNA precursor encoding plant genes. Alternatively, the second primer sequence may be based upon sequences derived from the cloning vector. For example, the skilled artisan can follow the RACE protocol (Frohman et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:8998-9002) to generate cDNAs by using PCR to amplify copies of the region between a single point in the transcript and the 3' or 5' end. Primers oriented in the 3' and 5' directions can be designed from the instant sequences. Using commercially available 3' RACE or 5' RACE systems (BRL), specific 3' or 5' cDNA fragments can be isolated (Ohara et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:5673-5677; Loh et al. (1989) *Science* 243:217-220). Products generated by the 3' and 5' RACE procedures can be combined to generate full-length cDNAs (Frohman and Martin (1989) *Techniques* 1:165). Consequently, a polynucleotide comprising a nucleotide sequence of at least one of 60 (preferably one of at least 40, most preferably one of at least 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, and 16 and the complement of such nucleotide sequences may be used in such methods to obtain a nucleic acid fragment encoding a substantial portion of an amino acid sequence of a polypeptide. The present invention relates to a method of obtaining a nucleic acid fragment encoding a substantial portion of a polypeptide of a gene (such as delta-6 desaturase or sphingolipid desaturase) preferably a



substantial portion of a plant polypeptide of a gene, comprising the steps of: synthesizing an oligonucleotide primer comprising a nucleotide sequence of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, and 16, and the complement of such nucleotide sequences; and amplifying a nucleic acid fragment (preferably a cDNA inserted in a cloning vector) using the oligonucleotide primer. The amplified nucleic acid fragment preferably will encode a portion of a polypeptide.

Availability of the instant nucleotide and deduced amino acid sequences facilitates immunological screening of cDNA expression libraries. Synthetic peptides representing portions of the instant amino acid sequences may be synthesized. These peptides can be used to immunize animals to produce polyclonal or monoclonal antibodies with specificity for peptides or proteins comprising the amino acid sequences. These antibodies can be then be used to screen cDNA expression libraries to isolate full-length cDNA clones of interest (Lerner (1984) *Adv. Immunol.* 36:1-34; Maniatis).

The nucleic acid fragments of the instant invention may be used to create transgenic plants in which the disclosed polypeptides are present at higher or lower levels than normal or in cell types or developmental stages in which they are not normally found. This would have the effect of altering the composition of fatty acids in those cells. Overexpression of delta-6 desaturase may allow the production of higher levels of tariric acid in *Picramnia*. Co-suppression of sphingolipid desaturase may allow for the production of higher levels of unsaturated corn and soybean oils.

Overexpression of the proteins of the instant invention may be accomplished by first constructing a chimeric gene in which the coding region is operably linked to a promoter capable of directing expression of a gene in the desired tissues at the desired stage of development. For reasons of convenience, the chimeric gene may comprise promoter sequences and translation leader sequences derived from the same genes. 3' Non-coding sequences encoding transcription termination signals may also be provided. The instant chimeric gene may also comprise one or more introns in order to facilitate gene expression.

Plasmid vectors comprising the instant chimeric gene can then be constructed. The choice of plasmid vector is dependent upon the method that will be used to transform host plants. The skilled artisan is well aware of the genetic elements that must be present on the plasmid vector in order to successfully transform, select and propagate host cells containing the chimeric gene. The skilled artisan will also recognize that different independent transformation events will result in different levels and patterns of expression (Jones et al. (1985) *EMBO J.* 4:2411-2418; De Almeida et al. (1989) *Mol. Gen. Genetics* 218:78-86), and thus that multiple events must be screened in order to obtain lines displaying the desired expression level and pattern. Such screening may be accomplished by Southern analysis of

DNA, Northern analysis of mRNA expression, Western analysis of protein expression, or phenotypic analysis.

For some applications it may be useful to direct the instant polypeptides to different cellular compartments, or to facilitate its secretion from the cell. It is thus envisioned that the chimeric gene described above may be further supplemented by altering the coding sequence to encode the instant polypeptides with appropriate intracellular targeting sequences such as transit sequences (Keegstra (1989) *Cell* 56:247-253), signal sequences or sequences encoding endoplasmic reticulum localization (Chrispeels (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53), or nuclear localization signals (Raikhel (1992) *Plant Phys.* 100:1627-1632) added and/or with targeting sequences that are already present removed. While the references cited give examples of each of these, the list is not exhaustive and more targeting signals of utility may be discovered in the future.

It may also be desirable to reduce or eliminate expression of genes encoding the instant polypeptides in plants for some applications. In order to accomplish this, a chimeric gene designed for co-suppression of the instant polypeptide can be constructed by linking a gene or gene fragment encoding that polypeptide to plant promoter sequences.

Alternatively, a chimeric gene designed to express antisense RNA for all or part of the instant nucleic acid fragment can be constructed by linking the gene or gene fragment in reverse orientation to plant promoter sequences. Either the co-suppression or antisense chimeric genes could be introduced into plants via transformation wherein expression of the corresponding endogenous genes are reduced or eliminated.

Molecular genetic solutions to the generation of plants with altered gene expression have a decided advantage over more traditional plant breeding approaches. Changes in plant phenotypes can be produced by specifically inhibiting expression of one or more genes by antisense inhibition or cosuppression (U.S. Patent Nos. 5,190,931, 5,107,065 and 5,283,323). An antisense or cosuppression construct would act as a dominant negative regulator of gene activity. While conventional mutations can yield negative regulation of gene activity these effects are most likely recessive. The dominant negative regulation available with a transgenic approach may be advantageous from a breeding perspective. In addition, the ability to restrict the expression of specific phenotype to the reproductive tissues of the plant by the use of tissue specific promoters may confer agronomic advantages relative to conventional mutations which may have an effect in all tissues in which a mutant gene is ordinarily expressed.

The person skilled in the art will know that special considerations are associated with the use of antisense or cosuppression technologies in order to reduce expression of particular genes. For example, the proper level of expression of sense or antisense genes may require the use of different chimeric genes utilizing different regulatory elements known to the skilled artisan. Once transgenic plants are obtained by one of the methods described above,

it will be necessary to screen individual transgenics for those that most effectively display the desired phenotype. The nature of these screens will generally be chosen on practical grounds, and is not an inherent part of the invention. For example, one can screen by looking for changes in gene expression by using antibodies specific for the protein encoded by the gene being suppressed, or one could establish assays that specifically measure enzyme activity. A preferred method will be one which allows large numbers of samples to be processed rapidly, since it will be expected that a large number of transformants will be negative for the desired phenotype.

The instant polypeptides (or portions thereof) may be produced in heterologous host cells, particularly in the cells of microbial hosts, and can be used to prepare antibodies to these proteins by methods well known to those skilled in the art. The antibodies are useful for detecting the polypeptides of the instant invention *in situ* in cells or *in vitro* in cell extracts. Preferred heterologous host cells for production of the instant polypeptides are microbial hosts. Microbial expression systems and expression vectors containing regulatory sequences that direct high level expression of foreign proteins are well known to those skilled in the art. Any of these could be used to construct a chimeric gene for production of the instant polypeptides. This chimeric gene could then be introduced into appropriate microorganisms via transformation to provide high level expression of the encoded delta-6 desaturase or sphingolipid desaturase. An example of a vector for high level expression of the instant polypeptides in a bacterial host is provided (Example 7).

All or a substantial portion of the nucleic acid fragments of the instant invention may also be used as probes for genetically and physically mapping the genes that they are a part of, and as markers for traits linked to those genes. Such information may be useful in plant breeding in order to develop lines with desired phenotypes. For example, the instant nucleic acid fragments may be used as restriction fragment length polymorphism (RFLP) markers. Southern blots (Maniatis) of restriction-digested plant genomic DNA may be probed with the nucleic acid fragments of the instant invention. The resulting banding patterns may then be subjected to genetic analyses using computer programs such as MapMaker (Lander et al. (1987) *Genomics* 1:174-181) in order to construct a genetic map. In addition, the nucleic acid fragments of the instant invention may be used to probe Southern blots containing restriction endonuclease-treated genomic DNAs of a set of individuals representing parent and progeny of a defined genetic cross. Segregation of the DNA polymorphisms is noted and used to calculate the position of the instant nucleic acid sequence in the genetic map previously obtained using this population (Botstein et al. (1980) *Am. J. Hum. Genet.* 32:314-331).

The production and use of plant gene-derived probes for use in genetic mapping is described in Bernatzky and Tanksley (1986) *Plant Mol. Biol. Reporter* 4:37-41. Numerous publications describe genetic mapping of specific cDNA clones using the methodology

outlined above or variations thereof. For example, F2 intercross populations, backcross populations, randomly mated populations, near isogenic lines, and other sets of individuals may be used for mapping. Such methodologies are well known to those skilled in the art.

Nucleic acid probes derived from the instant nucleic acid sequences may also be used for physical mapping (i.e., placement of sequences on physical maps; see Hoheisel et al. In: *Nonmammalian Genomic Analysis: A Practical Guide*, Academic press 1996, pp. 319-346, and references cited therein).

In another embodiment, nucleic acid probes derived from the instant nucleic acid sequences may be used in direct fluorescence *in situ* hybridization (FISH) mapping (Trask (1991) *Trends Genet.* 7:149-154). Although current methods of FISH mapping favor use of large clones (several to several hundred KB; see Laan et al. (1995) *Genome Res.* 5:13-20), improvements in sensitivity may allow performance of FISH mapping using shorter probes.

A variety of nucleic acid amplification-based methods of genetic and physical mapping may be carried out using the instant nucleic acid sequences. Examples include allele-specific amplification (Kazazian (1989) *J. Lab. Clin. Med.* 11:95-96), polymorphism of PCR-amplified fragments (CAPS; Sheffield et al. (1993) *Genomics* 16:325-332), allele-specific ligation (Landegren et al. (1988) *Science* 241:1077-1080), nucleotide extension reactions (Sokolov (1990) *Nucleic Acid Res.* 18:3671), Radiation Hybrid Mapping (Walter et al. (1997) *Nat. Genet.* 7:22-28) and Happy Mapping (Dear and Cook (1989) *Nucleic Acid Res.* 17:6795-6807). For these methods, the sequence of a nucleic acid fragment is used to design and produce primer pairs for use in the amplification reaction or in primer extension reactions. The design of such primers is well known to those skilled in the art. In methods employing PCR-based genetic mapping, it may be necessary to identify DNA sequence differences between the parents of the mapping cross in the region corresponding to the instant nucleic acid sequence. This, however, is generally not necessary for mapping methods.

Loss of function mutant phenotypes may be identified for the instant cDNA clones either by targeted gene disruption protocols or by identifying specific mutants for these genes contained in a maize population carrying mutations in all possible genes (Ballinger and Benzer (1989) *Proc. Natl. Acad. Sci USA* 86:9402-9406; Koes et al. (1995) *Proc. Natl. Acad. Sci USA* 92:8149-8153; Bensen et al. (1995) *Plant Cell* 7:75-84). The latter approach may be accomplished in two ways. First, short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols in conjunction with a mutation tag sequence primer on DNAs prepared from a population of plants in which Mutator transposons or some other mutation-causing DNA element has been introduced (see Bensen, *supra*). The amplification of a specific DNA fragment with these primers indicates the insertion of the mutation tag element in or near the plant gene encoding the instant polypeptides. Alternatively, the instant nucleic acid fragment may be used as a

hybridization probe against PCR amplification products generated from the mutation population using the mutation tag sequence primer in conjunction with an arbitrary genomic site primer, such as that for a restriction enzyme site-anchored synthetic adaptor. With either method, a plant containing a mutation in the endogenous gene encoding the instant polypeptides can be identified and obtained. This mutant plant can then be used to determine or confirm the natural function of the instant polypeptides disclosed herein.

### EXAMPLES

The present invention is further defined in the following Examples, in which all parts and percentages are by weight and degrees are Celsius, unless otherwise stated. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

#### EXAMPLE 1

##### Composition of cDNA Libraries; Isolation and Sequencing of cDNA Clones

cDNA libraries representing mRNAs from various Florida bitterbush, corn, soybean, and wheat tissues were prepared. The characteristics of the libraries are described below.

TABLE 2

cDNA Libraries from Florida Bitterbush, Corn, Soybean, and Wheat

Library	Tissue	Clone
pps	Developing Seeds of <i>Picramnia pentandra</i> (Florida bitterbush)	pps.pk0011.d5:fis
cde1c	Corn ( <i>Zea mays</i> , B13) Developing Embryo 20 DAP	cde1c.pk001.08:fis
sfl1	Soybean Immature Flower	sfl1.pk0031.d11
ssl	Soybean Seedling 5-10 Days After Germination	ssl.pk0017.b4:fis
wre1	Wheat Root From 7 Day Old Etiolated Seedling	wre1.pk0004.c7:fis

cDNA libraries may be prepared by any one of many methods available. For example, the cDNAs may be introduced into plasmid vectors by first preparing the cDNA libraries in Uni-ZAP™ XR vectors according to the manufacturer's protocol (Stratagene Cloning Systems, La Jolla, CA). The Uni-ZAP™ XR libraries are converted into plasmid libraries according to the protocol provided by Stratagene. Upon conversion, cDNA inserts will be contained in the plasmid vector pBluescript. In addition, the cDNAs may be introduced directly into pre-cut Bluescript II SK(+) vectors (Stratagene) using T4 DNA ligase (New England Biolabs), followed by transfection into DH10B cells according to the manufacturer's protocol (GIBCO BRL Products). Once the cDNA inserts are in plasmid

vectors, plasmid DNAs are prepared from randomly picked bacterial colonies containing recombinant pBluescript plasmids, or the insert cDNA sequences are amplified via polymerase chain reaction using primers specific for vector sequences flanking the inserted cDNA sequences. Amplified insert DNAs or plasmid DNAs are sequenced in dye-primer sequencing reactions to generate partial cDNA sequences (expressed sequence tags or “ESTs”; see Adams et al., (1991) *Science* 252:1651-1656). The resulting ESTs are analyzed using a Perkin Elmer Model 377 fluorescent sequencer.

## EXAMPLE 2

### Identification of cDNA Clones

cDNA clones encoding delta-6 desaturase or sphingolipid desaturase were identified by conducting BLAST (Basic Local Alignment Search Tool; Altschul et al. (1993) *J. Mol. Biol.* 215:403-410; see also [www.ncbi.nlm.nih.gov/BLAST/](http://www.ncbi.nlm.nih.gov/BLAST/)) searches for similarity to sequences contained in the BLAST “nr” database (comprising all non-redundant GenBank CDS translations, sequences derived from the 3-dimensional structure Brookhaven Protein Data Bank, the last major release of the SWISS-PROT protein sequence database, EMBL, and DDBJ databases). The cDNA sequences obtained in Example 1 were analyzed for similarity to all publicly available DNA sequences contained in the “nr” database using the BLASTN algorithm provided by the National Center for Biotechnology Information (NCBI). The DNA sequences were translated in all reading frames and compared for similarity to all publicly available protein sequences contained in the “nr” database using the BLASTX algorithm (Gish and States (1993) *Nat. Genet.* 3:266-272) provided by the NCBI. For convenience, the P-value (probability) of observing a match of a cDNA sequence to a sequence contained in the searched databases merely by chance as calculated by BLAST are reported herein as “pLog” values, which represent the negative of the logarithm of the reported P-value. Accordingly, the greater the pLog value, the greater the likelihood that the cDNA sequence and the BLAST “hit” represent homologous proteins.

## EXAMPLE 3

### Characterization of cDNA Clones Encoding Delta-6 Desaturase

The BLASTX search using the EST sequences from clones listed in Table 3 revealed similarity of the polypeptides encoded by the cDNAs to delta-6 desaturase from borage [*Borago officinalis*] (NCBI General Identifier No. 2062403). Shown in Table 3 are the BLAST results for the sequences of the entire cDNA inserts comprising the indicated cDNA clones (“FIS”):

**TABLE 3**  
BLAST Results for Sequences Encoding Polypeptides  
Homologous to Delta-6 Desaturase

Clone	Status	BLAST pLog Score 2062403
pps.pk0011.d5:fis	FIS	132.00

Sequence alignments and percent identity calculations were performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS* 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5. Sequence alignments and BLAST scores and probabilities indicate that the nucleic acid fragments comprising the instant cDNA clones encode a substantial portion of a delta-6 desaturase. These sequences represent the first *Picramnia pentandra* sequences encoding delta-6 desaturase.

#### EXAMPLE 4

##### Characterization of cDNA Clones Encoding Sphingolipid Desaturase

The BLASTX search using the EST sequences from clones listed in Table 4 revealed similarity of the polypeptides encoded by the cDNAs to sphingolipid desaturase from borage [*Borago officinalis*] (NCBI General Identifier No. 2062403), wheat [*Triticum aestivum*] (NCBI General Identifier No. 4104056), and sunflower [*Helianthus annuus*] (NCBI General Identifier No. 1040729). Shown in Table 4 are the BLAST results for the sequences of the entire cDNA inserts comprising the indicated cDNA clones ("FIS"), or contigs assembled from two or more ESTs ("Contig"):

**TABLE 4**  
BLAST Results for Sequences Encoding Polypeptides  
Homologous to Sphingolipid Desaturase

Clone	Status	NCBI General Identifier No.	BLAST pLog Score
ssl.pk0017.b4:fis	FIS	2062403	254.00
cde1c.pk001.o8:fis	FIS	4104056	254.00
wre1.pk0004.c7:fis	FIS	4104056	254.00
sfl1.pk0012.c5 sfl1.pk0031.d11	Contig	1040729	88.00

Figure 1 presents an alignment of the amino acid sequences set forth in SEQ ID NOs:2, 4, 6, 8, and 10 and the borage sequence (SEQ ID NO:11), wheat sequence (SEQ ID

NO:12), and sunflower sequence (SEQ ID NO:13). The data in Table 5 represents a calculation of the percent identity of the amino acid sequences set forth in SEQ ID NOs:2, 4, 6, 8, and 10, and the borage sequence (NCBI General Identifier No. 2062403, SEQ ID NO:11), wheat sequence (NCBI General Identifier No. 4104056, SEQ ID NO:12), and sunflower sequence (NCBI General Identifier No. 1040729, SEQ ID NO:13). The data in Table 5 represents a calculation of the percent identity of the amino acid sequence set forth in SEQ ID NOs:2, 4, 6, 8, and 10 and the delta 6 desaturase borage sequence (SEQ ID NO:11) and the wheat and sunflower sphingolipid desaturases (SEQ ID NOs:12 and 13, respectively).

TABLE 5

Percent Identity of Amino Acid Sequences Deduced From the Nucleotide Sequences of cDNA Clones Encoding Polypeptides Homologous to Delta-6 Desaturase and Sphingolipid Desaturase

SEQ ID NO.	Percent Identity to		
	2062403	4104056	1040729
2	54.0	49.2	51.4
8	62.7	55.2	70.3
4	54.0	79.5	53.7
10	54.0	99.8	53.5
6	49.8	48.6	53.0

The wheat sequence that is a virtual match to SEQ ID NO:10 was deposited into NCBI on January 29, 1999, almost two months after the priority date of the provisional filing of the present application. Therefore the wheat sequence that is claimed herein is the EST filed in the provisional application (herein labeled SEQ ID NO:17) that is comprised of amino acids 17-130 in SEQ ID NO:10.

Sequence alignments and percent identity calculations were performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5. Sequence alignments and BLAST scores and probabilities indicate that the nucleic acid fragments comprising the instant cDNA clones encode a substantial portion of a sphingolipid desaturase. These sequences represent the first monocot and first soybean sequences encoding sphingolipid desaturase.



### EXAMPLE 5

#### Expression of Chimeric Genes in Monocot Cells

A chimeric gene comprising a cDNA encoding the instant polypeptides in sense orientation with respect to the maize 27 kD zein promoter that is located 5' to the cDNA fragment, and the 10 kD zein 3' end that is located 3' to the cDNA fragment, can be constructed. The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites (NcoI or SmaI) can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the digested vector pML103 as described below.

Amplification is then performed in a standard PCR. The amplified DNA is then digested with restriction enzymes NcoI and SmaI and fractionated on an agarose gel. The appropriate band can be isolated from the gel and combined with a 4.9 kb NcoI-SmaI fragment of the plasmid pML103. Plasmid pML103 has been deposited under the terms of the Budapest Treaty at ATCC (American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209), and bears accession number ATCC 97366. The DNA segment from pML103 contains a 1.05 kb SalI-NcoI promoter fragment of the maize 27 kD zein gene and a 0.96 kb SmaI-SalI fragment from the 3' end of the maize 10 kD zein gene in the vector pGem9Zf(+) (Promega). Vector and insert DNA can be ligated at 15°C overnight, essentially as described (Maniatis). The ligated DNA may then be used to transform *E. coli* XL1-Blue (Epicurian Coli XL-1 Blue™; Stratagene). Bacterial transformants can be screened by restriction enzyme digestion of plasmid DNA and limited nucleotide sequence analysis using the dideoxy chain termination method (Sequenase™ DNA Sequencing Kit; U.S. Biochemical). The resulting plasmid construct would comprise a chimeric gene encoding, in the 5' to 3' direction, the maize 27 kD zein promoter, a cDNA fragment encoding the instant polypeptides, and the 10 kD zein 3' region.

The chimeric gene described above can then be introduced into corn cells by the following procedure. Immature corn embryos can be dissected from developing caryopses derived from crosses of the inbred corn lines H99 and LH132. The embryos are isolated 10 to 11 days after pollination when they are 1.0 to 1.5 mm long. The embryos are then placed with the axis-side facing down and in contact with agarose-solidified N6 medium (Chu et al. (1975) *Sci. Sin. Peking* 18:659-668). The embryos are kept in the dark at 27°C. Friable embryogenic callus consisting of undifferentiated masses of cells with somatic proembryoids and embryoids borne on suspensor structures proliferates from the scutellum of these immature embryos. The embryogenic callus isolated from the primary explant can be cultured on N6 medium and sub-cultured on this medium every 2 to 3 weeks.

The plasmid, p35S/Ac (obtained from Dr. Peter Eckes, Hoechst Ag, Frankfurt, Germany) may be used in transformation experiments in order to provide for a selectable marker. This plasmid contains the *Pat* gene (see European Patent Publication 0 242 236)

which encodes phosphinothricin acetyl transferase (PAT). The enzyme PAT confers resistance to herbicidal glutamine synthetase inhibitors such as phosphinothricin. The *pat* gene in p35S/Ac is under the control of the 35S promoter from Cauliflower Mosaic Virus (Odell et al. (1985) *Nature* 313:810-812) and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*.

The particle bombardment method (Klein et al. (1987) *Nature* 327:70-73) may be used to transfer genes to the callus culture cells. According to this method, gold particles (1  $\mu$ m in diameter) are coated with DNA using the following technique. Ten  $\mu$ g of plasmid DNAs are added to 50  $\mu$ L of a suspension of gold particles (60 mg per mL). Calcium chloride (50  $\mu$ L of a 2.5 M solution) and spermidine free base (20  $\mu$ L of a 1.0 M solution) are added to the particles. The suspension is vortexed during the addition of these solutions. After 10 minutes, the tubes are briefly centrifuged (5 sec at 15,000 rpm) and the supernatant removed. The particles are resuspended in 200  $\mu$ L of absolute ethanol, centrifuged again and the supernatant removed. The ethanol rinse is performed again and the particles resuspended in a final volume of 30  $\mu$ L of ethanol. An aliquot (5  $\mu$ L) of the DNA-coated gold particles can be placed in the center of a Kapton™ flying disc (Bio-Rad Labs). The particles are then accelerated into the corn tissue with a Biolistic™ PDS-1000/He (Bio-Rad Instruments, Hercules CA), using a helium pressure of 1000 psi, a gap distance of 0.5 cm and a flying distance of 1.0 cm.

For bombardment, the embryogenic tissue is placed on filter paper over agarose-solidified N6 medium. The tissue is arranged as a thin lawn and covered a circular area of about 5 cm in diameter. The petri dish containing the tissue can be placed in the chamber of the PDS-1000/He approximately 8 cm from the stopping screen. The air in the chamber is then evacuated to a vacuum of 28 inches of Hg. The macrocarrier is accelerated with a helium shock wave using a rupture membrane that bursts when the He pressure in the shock tube reaches 1000 psi.

Seven days after bombardment the tissue can be transferred to N6 medium that contains glufosinate (2 mg per liter) and lacks casein or proline. The tissue continues to grow slowly on this medium. After an additional 2 weeks the tissue can be transferred to fresh N6 medium containing glufosinate. After 6 weeks, areas of about 1 cm in diameter of actively growing callus can be identified on some of the plates containing the glufosinate-supplemented medium. These calli may continue to grow when sub-cultured on the selective medium.

Plants can be regenerated from the transgenic callus by first transferring clusters of tissue to N6 medium supplemented with 0.2 mg per liter of 2,4-D. After two weeks the tissue can be transferred to regeneration medium (Fromm et al. (1990) *Bio/Technology* 8:833-839).

## EXAMPLE 6

### Expression of Chimeric Genes in Dicot Cells

A seed-specific expression cassette composed of the promoter and transcription terminator from the gene encoding the  $\beta$  subunit of the seed storage protein phaseolin from the bean *Phaseolus vulgaris* (Doyle et al. (1986) *J. Biol. Chem.* 261:9228-9238) can be used for expression of the instant polypeptides in transformed soybean. The phaseolin cassette includes about 500 nucleotides upstream (5') from the translation initiation codon and about 1650 nucleotides downstream (3') from the translation stop codon of phaseolin. Between the 5' and 3' regions are the unique restriction endonuclease sites Nco I (which includes the ATG translation initiation codon), Sma I, Kpn I and Xba I. The entire cassette is flanked by Hind III sites.

The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the expression vector. Amplification is then performed as described above, and the isolated fragment is inserted into a pUC18 vector carrying the seed expression cassette.

Soybean embryos may then be transformed with the expression vector comprising sequences encoding the instant polypeptides. To induce somatic embryos, cotyledons, 3-5 mm in length dissected from surface sterilized, immature seeds of the soybean cultivar A2872, can be cultured in the light or dark at 26°C on an appropriate agar medium for 6-10 weeks. Somatic embryos which produce secondary embryos are then excised and placed into a suitable liquid medium. After repeated selection for clusters of somatic embryos which multiplied as early, globular staged embryos, the suspensions are maintained as described below.

Soybean embryogenic suspension cultures can be maintained in 35 mL liquid media on a rotary shaker, 150 rpm, at 26°C with florescent lights on a 16:8 hour day/night schedule. Cultures are subcultured every two weeks by inoculating approximately 35 mg of tissue into 35 mL of liquid medium.

Soybean embryogenic suspension cultures may then be transformed by the method of particle gun bombardment (Klein et al. (1987) *Nature* (London) 327:70-73, U.S. Patent No. 4,945,050). A DuPont Biolistic™ PDS1000/HE instrument (helium retrofit) can be used for these transformations.

A selectable marker gene which can be used to facilitate soybean transformation is a chimeric gene composed of the 35S promoter from Cauliflower Mosaic Virus (Odell et al. (1985) *Nature* 313:810-812), the hygromycin phosphotransferase gene from plasmid pJR225 (from *E. coli*; Gritz et al. (1983) *Gene* 25:179-188) and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*. The seed expression

cassette comprising the phaseolin 5' region, the fragment encoding the instant polypeptides and the phaseolin 3' region can be isolated as a restriction fragment. This fragment can then be inserted into a unique restriction site of the vector carrying the marker gene.

To 50  $\mu$ L of a 60 mg/mL 1  $\mu$ m gold particle suspension is added (in order): 5  $\mu$ L DNA (1  $\mu$ g/ $\mu$ L), 20  $\mu$ l spermidine (0.1 M), and 50  $\mu$ L CaCl<sub>2</sub> (2.5 M). The particle preparation is then agitated for three minutes, spun in a microfuge for 10 seconds and the supernatant removed. The DNA-coated particles are then washed once in 400  $\mu$ L 70% ethanol and resuspended in 40  $\mu$ L of anhydrous ethanol. The DNA/particle suspension can be sonicated three times for one second each. Five  $\mu$ L of the DNA-coated gold particles are then loaded on each macro carrier disk.

Approximately 300-400 mg of a two-week-old suspension culture is placed in an empty 60x15 mm petri dish and the residual liquid removed from the tissue with a pipette. For each transformation experiment, approximately 5-10 plates of tissue are normally bombarded. Membrane rupture pressure is set at 1100 psi and the chamber is evacuated to a vacuum of 28 inches mercury. The tissue is placed approximately 3.5 inches away from the retaining screen and bombarded three times. Following bombardment, the tissue can be divided in half and placed back into liquid and cultured as described above.

Five to seven days post bombardment, the liquid media may be exchanged with fresh media, and eleven to twelve days post bombardment with fresh media containing 50 mg/mL hygromycin. This selective media can be refreshed weekly. Seven to eight weeks post bombardment, green, transformed tissue may be observed growing from untransformed, necrotic embryogenic clusters. Isolated green tissue is removed and inoculated into individual flasks to generate new, clonally propagated, transformed embryogenic suspension cultures. Each new line may be treated as an independent transformation event. These suspensions can then be subcultured and maintained as clusters of immature embryos or regenerated into whole plants by maturation and germination of individual somatic embryos.

#### EXAMPLE 7

##### Expression of Chimeric Genes in Microbial Cells

The cDNAs encoding the instant polypeptides can be inserted into the T7 *E. coli* expression vector pBT430. This vector is a derivative of pET-3a (Rosenberg et al. (1987) *Gene* 56:125-135) which employs the bacteriophage T7 RNA polymerase/T7 promoter system. Plasmid pBT430 was constructed by first destroying the EcoR I and Hind III sites in pET-3a at their original positions. An oligonucleotide adaptor containing EcoR I and Hind III sites was inserted at the BamH I site of pET-3a. This created pET-3aM with additional unique cloning sites for insertion of genes into the expression vector. Then, the Nde I site at the position of translation initiation was converted to an Nco I site using oligonucleotide-directed mutagenesis. The DNA sequence of pET-3aM in this region, 5'-CATATGG, was converted to 5'-CCCATGG in pBT430.

Plasmid DNA containing a cDNA may be appropriately digested to release a nucleic acid fragment encoding the protein. This fragment may then be purified on a 1% NuSieve GTG™ low melting agarose gel (FMC). Buffer and agarose contain 10 µg/ml ethidium bromide for visualization of the DNA fragment. The fragment can then be purified from the agarose gel by digestion with GELase™ (Epicentre Technologies) according to the manufacturer's instructions, ethanol precipitated, dried and resuspended in 20 µL of water. Appropriate oligonucleotide adapters may be ligated to the fragment using T4 DNA ligase (New England Biolabs, Beverly, MA). The fragment containing the ligated adapters can be purified from the excess adapters using low melting agarose as described above. The vector pBT430 is digested, dephosphorylated with alkaline phosphatase (NEB) and deproteinized with phenol/chloroform as described above. The prepared vector pBT430 and fragment can then be ligated at 16°C for 15 hours followed by transformation into DH5 electrocompetent cells (GIBCO BRL). Transformants can be selected on agar plates containing LB media and 100 µg/mL ampicillin. Transformants containing the gene encoding the instant polypeptides are then screened for the correct orientation with respect to the T7 promoter by restriction enzyme analysis.

For high level expression, a plasmid clone with the cDNA insert in the correct orientation relative to the T7 promoter can be transformed into *E. coli* strain BL21(DE3) (Studier et al. (1986) *J. Mol. Biol.* 189:113-130). Cultures are grown in LB medium containing ampicillin (100 mg/L) at 25°C. At an optical density at 600 nm of approximately 1, IPTG (isopropylthio-β-galactoside, the inducer) can be added to a final concentration of 0.4 mM and incubation can be continued for 3 h at 25°. Cells are then harvested by centrifugation and re-suspended in 50 µL of 50 mM Tris-HCl at pH 8.0 containing 0.1 mM DTT and 0.2 mM phenyl methylsulfonyl fluoride. A small amount of 1 mm glass beads can be added and the mixture sonicated 3 times for about 5 seconds each time with a microprobe sonicator. The mixture is centrifuged and the protein concentration of the supernatant determined. One µg of protein from the soluble fraction of the culture can be separated by SDS-polyacrylamide gel electrophoresis. Gels can be observed for protein bands migrating at the expected molecular weight.

#### EXAMPLE 8

##### Expression of *Picramnia* EST pps.pk0011.d5 in Somatic Soybean Embryos

As one method of assaying the function of the polypeptide encoded by the *Picramnia* EST pps.pk0011.d5, the corresponding cDNA was expressed in somatic soybean embryos, and the fatty acid composition of the resulting transgenic tissue was examined for alterations accompanying the expression of the cDNA.

A plasmid pZBL100 containing chimeric genes to allow expression of hygromycin B phosphotransferase in certain bacteria and in plant cells was constructed from the following genetic elements: (a) T7 promoter + Shine-Delgarno / hygromycin B

phosphotransferase (HPT) / T7 terminator sequence, (b) 35S promoter from cauliflower mosaic virus (CaMV) / hygromycin B phosphotransferase (HPT) / nopaline synthase (NOS3' from *Agrobacterium tumefaciens* T-DNA, and (c) pSP72 plasmid vector [from Promega] with  $\beta$ -lactamase coding region (ampicillin resistance gene) removed.

5 The hygromycin B phosphotransferase gene was amplified by PCR from *E. coli* strain W677, which contained a *Klebsiella* derived plasmid pJR225. Starting with the pSP72 vector the elements were assembled into a single plasmid using standard cloning methods (Maniatis).

10 Plasmid pZBL100 thus contains the T7 promoter/HPT/T7 terminator cassette for expression of the HPT enzyme in certain strains of *E. coli*, such as NovaBlue (DE3) (Novagen), that are lysogenic for lambda DE3 (which carries the T7 RNA Polymerase gene under *lacUV5* control). Plasmid pZBL100 also contains the 35S/HPT/NOS cassette for constitutive expression of the HPT enzyme in plants, such as soybean. These two  
15 expression systems allow selection for growth in the presence of hygromycin to be used as a means of identifying cells that contain the plasmid in both bacterial and plant systems.

PZBL100 also contains three unique restriction endonuclease sites suitable for the cloning of other chimeric genes into this vector.

A plasmid for expression of the cDNA encoding fatty acid modifying enzymes under the control of the soybean  $\beta$ -conglycinin promoter (Beachy et al., (1985) EMBO J. 4:3047-3053) can be constructed. The construction of this vector was facilitated by the use  
20 of plasmids pCW109 and pML18, both of which have been described (see World Patent Publication No. WO 94/11516).

A unique *NotI* site was introduced into the cloning region between the  $\beta$ -conglycinin promoter and the phaseolin 3' end in pCW109 by digestion with *NcoI* and  
25 *XbaI* followed by removal of the single stranded DNA ends with mung bean exonuclease. *NotI* linkers (New England Biochemical catalog number NEB 1125) were ligated into the linearized plasmid to produce plasmid pAW35. The single *NotI* site in pML18 was destroyed by digestion with *NotI*, filling in the single stranded ends with dNTPs and Klenow fragment followed by re-ligation of the linearized plasmid. The modified pML18  
30 was then digested with *HindIII* and treated with calf intestinal phosphatase.

The  $\beta$ -conglycinin:*NotI*:phaseolin expression cassette in pAW35 was removed by digestion with *HindIII* and the 1.8 kB fragment was isolated by agarose gel electrophoresis. The isolated fragment was ligated into the modified and linearized pML18 construction described above. A clone with the desired orientation was identified by  
35 digestion with *NotI* and *XbaI* to release a 1.08 kB fragment indicating that the orientation of the  $\beta$ -conglycinin transcription unit was the same as the selectable marker transcription unit. The resulting plasmid was given the name pBS19

*Hind*III is one of the unique cloning sites available in pZBL100. To assemble the final expression cassette, pBS19 and pZBL100 were both digested with *Hind*III. The  $\beta$ -conglycinin containing fragment from pBS19 was isolated by gel electrophoresis and ligated into the digested pZBL100, which had been treated with calf alkaline phosphatase.

5 The resulting plasmid was named pKS67.

The cDNA insert for the *Picramnia* EST pk0011.d5 was amplified by PCR to generate flanking *Not*I sites to allow for cloning into the corresponding restriction site of plasmid pKS67. The 5' and 3' oligonucleotide primers used for the amplification of the *Picramnia* cDNA are provided in SEQ ID NO:14 and 15, respectively.

10 PCR reactions were conducted using *Pfu* polymerase (Stratagene) and the cDNA corresponding to EST pps.pk0011.d5 as the template. Products from amplification reactions were purified and subcloned into pCR-Script Amp SK(+). The PCR product was then moved as a *Not*I fragment into the soybean expression vector pKS67.

15 Gene fusions of the *Picramnia* cDNA with the conglycinin promoter and phaseolin termination sequences in vector pKS67 were introduced into soybean embryos using the particle bombardment method of transformation. To induce somatic embryos, cotyledons, 3-5 mm in length dissected from surface sterilized, immature seeds of a soybean cultivar, such as A2872, can be cultured in the light or dark at 26°C on an appropriate agar medium for 6-10 weeks. Somatic embryos that produce secondary embryos were then excised and  
20 placed into a suitable liquid medium. After repeated selection for clusters of somatic embryos that multiplied as early, globular staged embryos, the suspensions were maintained as described below.

Soybean embryogenic suspension cultures were maintained in 35 mL liquid media on a rotary shaker, 150 rpm, at 26°C with fluorescent lights on a 16:8 hour day/night schedule.  
25 Cultures were subcultured every two weeks by inoculating approximately 35 mg of tissue into 35 mL of liquid medium.

Soybean embryogenic suspension cultures were then transformed with the vector pKS67 containing the *Picramnia* cDNA for EST pps.pk0011.d5 by the method of particle gun bombardment (Klein et al. (1987) *Nature* (London) 327:70, U.S. Patent No. 4,945,050).  
30 A DuPont Biolistic PDS1000/HE instrument (helium retrofit) was used for these transformations.

To 50 mL of a 60 mg/mL 1 mm gold particle suspension were added (in order): 5 mL DNA (1 mg/mL), 20 ml spermidine (0.1 M), and 50 mL  $\text{CaCl}_2$  (2.5 M). The particle preparation was then agitated for three minutes, spun in a microfuge for 10 seconds and the  
35 supernatant removed. The DNA-coated particles were then washed once in 400 mL 70% ethanol and resuspended in 40 mL of anhydrous ethanol. The DNA/particle suspension was sonicated three times for one second each. Five mL of the DNA-coated gold particles was then loaded on each macro carrier disk.

Approximately 300-400 mg of a two-week-old suspension culture was placed in an empty 60x15-mm petri dish and the residual liquid removed from the tissue with a pipette. For each transformation experiment, approximately 5 to 10 plates of tissue were bombarded. Membrane rupture pressure was set at 1100 psi and the chamber was evacuated to a vacuum of 28 inches mercury. The tissue was placed approximately 3.5 inches away from the retaining screen and bombarded three times. Following bombardment, the tissue was divided in half and placed back into liquid and cultured as described above.

Five to seven days post bombardment, the liquid media was exchanged with fresh media, and eleven to twelve days post bombardment with fresh media containing 50 mg/mL hygromycin. This selective media was refreshed weekly. Seven to eight weeks post bombardment, green, transformed tissue was observed growing from untransformed, necrotic embryogenic clusters. Isolated green tissue was removed and inoculated into individual flasks to generate new, clonally propagated, transformed embryogenic suspension cultures. Each new line was treated as an independent transformation event. These suspensions were then subcultured and maintained as clusters of immature embryos.

Transgenic soybean embryos selected and maintained in this manner were analyzed for alterations in fatty acid composition. Individual embryos expressing the cDNA for the *Picramnia* EST pps.pk0011.d5 were homogenized in 1% (w/v) sodium methoxide in methanol. Fatty acid methyl esters resulting from this transesterification step were analyzed by both GC and GC-MS as described elsewhere (Hitz et al. (1994) Plant Physiol. 105:635-641). Using this methodology, individual somatic soybean embryos from transformation event MS151-5-4 were found to contain a fatty acid whose methyl ester had a retention time and mass spectrum consistent with that of gamma-linolenic acid (18:3 $\Delta^{6,9,12}$ ). This fatty acid is not normally present in somatic soybean embryos. In individual embryos expressing the cDNA for *Picramnia* EST pps.pk0011.d5, gamma-linolenic acid was detected in amounts as high as 1.2% (wt/wt) of the total fatty acids. Accompanying the occurrence of gamma-linolenic acid in transgenic somatic soybean embryos was an unidentified compound with a retention time intermediate to that of methyl  $\alpha$ -linolenic acid (18:3 $\Delta^{9,12,15}$ ) and methyl eicosanoic acid (20:0) on a polar GC column, such as Omegawax 320 (Supelco). Analysis by GC-MS indicated that this compound had an apparent molecular ion of 294 *m/z*. This compound accounted for ...0.5% of the total fatty acids of individual embryos from transformation event MS151-5-4 and was only found in embryos that contained detectable amounts of gamma-linolenic acid. Overall, the presence of gamma-linolenic acid in transgenic somatic soybean embryos confirms that the introduced gene is capable of enhancing a  $\Delta^6$  desaturation event in a tissue that normally does not accumulate fatty acids with these bonds. The production of tariric acid by this enzyme cannot be addressed due to the low overall accumulation of  $\Delta^6$  containing fatty acids. Isolation of embryos expressing higher levels of the introduced gene may resolve this question.



Various modifications of the invention in addition to those shown and described herein will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

5 The disclosure of each reference set forth above is incorporated herein by reference in its entirety.

## CLAIMS

What is claimed is:

1. An isolated polynucleotide comprising a first nucleotide sequence encoding a polypeptide of at least 60 amino acids that has at least 55% identity based on the Clustal method of alignment when compared to a delta-6 desaturase polypeptide of SEQ ID NO:2, a second nucleotide sequence encoding a first polypeptide of at least 114 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of a sphingolipid desaturase polypeptide of SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, and SEQ ID NO:17 or a third nucleotide sequence comprising the complement of the first or second nucleotide sequences.
2. The isolated polynucleotide of Claim 1, wherein the isolated polynucleotide consists of a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, and 16 that codes for the polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8, and 17.
3. The isolated polynucleotide of Claim 1 wherein the nucleotide sequences are DNA.
4. The isolated polynucleotide of Claim 1 wherein the nucleotide sequences are RNA.
5. A chimeric gene comprising the isolated polynucleotide of Claim 1 operably linked to suitable regulatory sequences.
6. An isolated host cell comprising the chimeric gene of Claim 5.
7. An isolated host cell comprising an isolated polynucleotide of Claim 1 or Claim 3.
8. The isolated host cell of Claim 7 wherein the isolated host is selected from the group consisting of yeast, bacteria, plant, and virus.
9. A virus comprising the isolated polynucleotide of Claim 1.
10. A polypeptide of at least 60 amino acids that has at least 55% identity based on the Clustal method of alignment when compared to the polypeptide of SEQ ID NO:2, or at least 114 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, and SEQ ID NO:17.
11. A method of selecting an isolated polynucleotide that affects the level of expression of a desaturase polypeptide in a plant cell, the method comprising the steps of:
  - (a) constructing an isolated polynucleotide comprising a nucleotide sequence of at least one of 30 contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs: 1, 3, 5, 7, and 16 and the complement of such nucleotide sequences;
  - (b) introducing the isolated polynucleotide into a plant cell;

(c) measuring the level of a polypeptide in the plant cell containing the polynucleotide; and

(d) comparing the level of polypeptide in the plant cell containing the isolated polynucleotide with the level of polypeptide in a plant cell that does not contain the isolated polynucleotide.

12. The method of Claim 11 wherein the isolated polynucleotide consists of a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, and 16 that codes for the polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8, and 17.

13. A method of selecting an isolated polynucleotide that affects the level of expression of a delta-6 desaturase or sphingolipid desaturase polypeptide in a plant cell, the method comprising the steps of:

(a) constructing an isolated polynucleotide of Claim 1;

(b) introducing the isolated polynucleotide into a plant cell;

(c) measuring the level of polypeptide in the plant cell containing the polynucleotide; and

(d) comparing the level of polypeptide in the plant cell containing the isolated polynucleotide with the level of polypeptide in a plant cell that does not contain the polynucleotide.

14. A method of obtaining a nucleic acid fragment encoding a delta-6 desaturase or sphingolipid desaturase polypeptide comprising the steps of:

(a) synthesizing an oligonucleotide primer comprising a nucleotide sequence of at least one of 40 contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, and 16 and the complement of such nucleotide sequences; and

(b) amplifying a nucleic acid sequence using the oligonucleotide primer.

15. A method of obtaining a nucleic acid fragment encoding the amino acid sequence encoding a delta-6 desaturase or sphingolipid desaturase polypeptide comprising the steps of:

(a) probing a cDNA or genomic library with an isolated polynucleotide comprising a nucleotide sequence of at least one of 30 contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, and 16 and the complement of such nucleotide sequences;

(b) identifying a DNA clone that hybridizes with the isolated polynucleotide;

(c) isolating the identified DNA clone; and

(d) sequencing the cDNA or genomic fragment that comprises the isolated DNA clone.

16. A method for positive selection of a transformed cell comprising:

(a) transforming a host cell with the chimeric gene of Claim 5; and

(b) growing the transformed host cell under conditions suitable for the expression of the chimeric gene allowing expression of the polynucleotide in an amount to alter the concentration of fatty acids with delta-6 double bonds in the host cell to provide a positive selection means.

5

17. The method of Claim 16 wherein the host cell is selected from the group consisting of plant cells and procaryotes.

18. The method of Claim 16 wherein levels of tariric acid are altered.

19. A method for positive selection of a transformed cell comprising:

10

(a) transforming a plant cell with the chimeric gene of Claim 5;

(b) growing a plant from the transformed plant cell of step (a) allowing expression of the polynucleotide in an amount to alter the concentration of fatty acids with delta-6 double bonds in the seeds of the plant to provide a positive selection means.

FIGURE 1

bitterbush_[SIN_2]	MEEPK-----KHISQADLAKHKQPGDLWISIKGKVYDISKW
soybean_[SIN_8]	M-----EVVEKEK-----KYITSEELKGNKEGDLWISIQGKVYNVSDW
borage_[gi_2062403]	MA-----A-----QIK-----KYITSDELKNHDKPGDLWISIQGKAYDVSDW
corn_[SIN_4]	MP-----PSVDAMPAPGDAAGADVVRMISSELRAHASADDLWISISGDVYDVTPW
wheat_[SIN_10]	MARTGLADATAPEADAMPAASK--DAADVVRMISTKELQAHAAADDLWISISGDVYDVTPW
wheat_[gi_4104056]	MARTGLADATAPEADAMPAASK--DAADVVRMISTKELQAHAAADDLWISISGDVYDVTPW
soybean_[SIN_6]	LP-----A-----
sunflower_[gi_1040729]	MV-----SPSIEVLNSIADGK-----KYITSKELKKNPNPDLWISILGKVYNVTEW
bitterbush_[SIN_2]	TKEHPGGELPLLSFAGQDVTDAFIAYHPGTAWQYLD RFFTYGVQDYVSVSEMSKDYRRLV
soybean_[SIN_8]	VKEHPGGDVPIISNLAGQDVTDAFIAYHPGTAWSHLEKFFTYGHLSDFKVSEVSKDYRKLA
borage_[gi_2062403]	VKDHPGGSFPLKSLAGQEVTDAFVAFHPASTWKNL DKFFTYGHLSDFKVSEVSKDYRKLV
corn_[SIN_4]	LPHPGGDLPLLTLAGQDATDAFAAYHPPSARPLLR RFFVG-RLSDYAVSPASADYRRL
wheat_[SIN_10]	LRHHPGGEVPLITLAGQDATDAFMAYHPPSVRPLLR RFFVG-RLSDYTVPPASADFRRL
wheat_[gi_4104056]	LRHHPGGEVPLITLAGQDATDAFMAYHPPSVRPLLR RFFVG-RLTDYTVPPASADFRRL
soybean_[SIN_6]	-----FSTSHRLSDHTVSAASSDYRKLF
sunflower_[gi_1040729]	AKEHPGGDAPLINLAGQDVTDAFIAFHPGTAWKHL DKLFTGYHLKDYQVSDISRDKLA
bitterbush_[SIN_2]	SEFSKMGLFKTPGKGVYCSIFFVSVLFALSVYGVLYCKSTWAHLCSGLLMGMLWLQSGWV
soybean_[SIN_8]	SEFSKLGLEFDTKGHVTSCTLASVAVMFLIVLYGVLRCTSVWAHLGSGMLLGLLWMQSA YV
borage_[gi_2062403]	FEFSKMGLYDKKGHIMFATLCFIAMLFAMSVYGVLFCEGVLVHLFSGCLMGFLWIQSGWI
corn_[SIN_4]	AQLSSAGLFEVGPVKVQLVLMVLFYAALYVLACASAWAHLLAGGLIGFVWIQSGWM
wheat_[SIN_10]	AQLSSAGLFEVGHTPKFLLVAMSVLFCIALYCVLACSTGAHMFAGGLIGFVWIQSGWI
wheat_[gi_4104056]	AQLSSAGLFEVGHTPKFLLVAMSVLFCIALYCVLACSTGAHMFAGGLIGFVWIQSGWI
soybean_[SIN_6]	SDLSALNLFNRKGHTTISLLSLITLPLSVCGVLFSDSTFVHVLAAALIGFLWIQSGWI
sunflower_[gi_1040729]	SEFAKAGMFEKKGHGVIYSLCFVSLLLSACVYGVLYSGSFWIHMLSGAILGLAWMQIAYL

FIGURE 1

bitterbush_[SIN_2]	GHDSCHYQVMPNKRKLRLFQIIAGNVIAGSVAVWVKLDHNTTHHFACNSANLDPDIQHLPI
soybean_[SIN_8]	GHDSGHYVVMTTNGFNKVAQILSGNCLTGISIAWKKWTHNAHHIACNSLDHDPDLQHMPV
borage_[gi_2062403]	GHDAAGHYMVVSDSRNLKFMGIFAANCLSGISIGWKKWNHNAHHIACNSLEYDQYIPF
corn_[SIN_4]	GHDSGHHRIITGHPVLDVVQVLSGNCLTGLSIAWKKCNHNTTHHIAACNSLDHDPDLQHMPV
wheat_[SIN_10]	GHDSGHHQITRHPALNRLQVVSGNCLTGLGIAWKKFNHNTTHHISCNSLDHDPDLQHLPV
wheat_[gi_4104056]	GHDSGHHQITRHPALNRLQVVSGNCLTGLGIAWKKFNHNTTHHISCNSLDHDPDLQHLPV
soybean_[SIN_6]	GHDSGHYNVMLSRRLNRAIQILSGNLAGISIGWKKWNHNAHHIACNSLDYDQYIPF
sunflower_[gi_1040729]	GHDAAGHYQMMATRGNKFAGIFIGNCITGISIAWKKWTHNAHHIACNSLDYDQYIPF
bitterbush_[SIN_2]	IAISPKFFNSLTSYHNCMTYDRAARFFVSFQHWTFYPALLSVRLYLFIKSVVFSNN
soybean_[SIN_8]	FAVSSRFFNSITSHFYGRKLEDFIAREFLICYQHFTFYPVCMCVARNLYLQITILLFSR-
borage_[gi_2062403]	LVVSSKFFGSLTSHFYEKRLTFDSLSRFFVSQHWTFYPIMCAARLNMYVQSLIMLLTK-
corn_[SIN_4]	FAVSPKLFNGIWSYFYQRTLAFDAASKFFISYQHWTFYPVVMCIARINLLAQSAFLVLT-
wheat_[SIN_10]	FAVSTKLFNNLWSVCYERTLAFDAISKFFVSQHWTFYPVVMGEARINLLVQSIIVFLITQ-
wheat_[gi_4104056]	FAVSTKLFNNLWSVCYERTLAFDAISKFFVSQHWTFYPVVMGEARINLLVQSIIVFLITQ-
soybean_[SIN_6]	FAVSSRFFNSITSHXYGRKXEFDXIAXFLICYQHFTFYPVCMCVARNLYLQITILLFSR-
sunflower_[gi_1040729]	LAVSSKLFNSITSVFYGRQLTFDPLARFFVSQHYLYYPIMCVARNLYLQITILLISK-
bitterbush_[SIN_2]	KRVYKRSQEILGYAAFLTMYSLLSRLPNWPERVMYFTSCLAVAGFQHWQFSLNHFASNV
soybean_[SIN_8]	RKVQDRALNIMGILVFWTWFPPLLVSCLPNWPERVMFVLASFVCSIQHIQFCLNHFANV
borage_[gi_2062403]	RNVSYRAHELGLCLVFSIWYPLLVSCLPNWGERIMFVIAASLSVTGMQVQFSLNHFSSSV
corn_[SIN_4]	KRVPQRLLEIAGVATFWAWYPLLVASLPNWWERVAFVLFSFTICGIQHVQFCLNHFSSDV
wheat_[SIN_10]	KKVRQRLWEIAGVAAFVWYPLLVSCLPNWWERVAFVLASFVITGIGHVQFCLNHFSSAV
wheat_[gi_4104056]	KKVRQRLWEIAGVAAFVWYPLLVSCLPNWWERVAFVLASFVITGIGHVQFCLNHFSSAV
soybean_[SIN_6]	XKVQDRALNIMGILVFWTWFLFLA-----LLFV-----PIQHIQFWLNHLAENL
sunflower_[gi_1040729]	RKIPDRGLNILGTLIFWTWFPPLLVSRLPNWPERVAFVLVSFCVTGIGIHIQFTLNHFSGDV

FIGURE 1

bitterbush_[SIN_2]	YTGLPSGNDWFHQQTGTLNITASAWNDWFHGGGLHFQIEHHLFPRMPKCHFRKISPIVNK
soybean_[SIN_8]	YVGPPSGNDWFEEKQTSGLDISCASSMDWFFGGLQFQLEHHLFPRLPRLPRCQLRKISPLVSD
borage_[gi_2062403]	YVGKPKGNWFEKQTDGTLDISCPPWMDWFHGGGLQFQIEHHLFPRMPRCNLRKISPYVIE
corn_[SIN_4]	YVGPPKGNWFEKQTAGTLILCSPWMDWFHGGGLQFQIEHHLFPRLPRLPRCHLRKVAPVRD
wheat_[SIN_10]	YVGPPKGNWFERQTAGTLIDIKSPWMDWFHGGGLQFQVEHHLFPRLPRLPRCHYRMVAPIVRD
wheat_[gi_4104056]	YVGPPKGNWFERQTAGTLIDIKSPWMDWFHGGGLQFQVEHHLFPRLPRLPRCHYRMVAPIVRD
soybean_[SIN_6]	YXG-----
sunflower_[gi_1040729]	YVGPPKGNWFEKQTRGTIDACSSWMDWFFGGLQFQLEHHLFPRLPRLCHLRSISPICRE
bitterbush_[SIN_2]	LCQKHNLSEYETATMWEANKMVYSTLRVAMEAKDVTK-PVPKNMVWEAMNTEFG
soybean_[SIN_8]	LCKKHNLPSRSLSFWEANQWTIRTLRTAALQARDLTN-PAPKNLLWEAVNTHG
borage_[gi_2062403]	LCKKHNLPPYNYASFESKANEMTLRTLRLNTALQARDITK-PLPKNLVWEALHTHG
corn_[SIN_4]	LCKKHGLTYSAAATFWGANVLTWKTLRAAALQARTATSGGAPKNLVWEAVNTHG
wheat_[SIN_10]	LCKKHGLSYGAATFWEANVMTWKTLRAAALQAREATTGAAPKNLVWEALNTHG
wheat_[gi_4104056]	LCKKHGLSYGAATFWEANVMTWKTLRAAALQAREATTGAAPKNLVWEALNTHG
soybean_[SIN_6]	-----
sunflower_[gi_1040729]	LCKKYNLPYVSLSFYDANVTTLKTLRTAALQARDLTN-PAPQNLAWAEAFNTHG

SEQUENCE LISTING

&lt;110&gt; E. I. du Pont de Nemours and Company

&lt;120&gt; Membrane-Bound Desaturases

&lt;130&gt; BB1264

&lt;140&gt;

&lt;141&gt;

&lt;150&gt; 60/110,784

&lt;151&gt; 1998-December-03

&lt;160&gt; 17

&lt;170&gt; Microsoft Office 97

&lt;210&gt; 1

&lt;211&gt; 1471

&lt;212&gt; DNA

&lt;213&gt; Picramnia pentandra

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (1402)

&lt;400&gt; 1

```

cttccttggt cctggaattt tcaaatcact tcctctgttg cacttcaatg gaagagccaa 60
agaagcacat ttcgcaagca gaccttgcaa agcataagca accaggagat ttatggatct 120
ctatcaaggg aaaagtttac gatatctcca agtggaactaa agagcatccc ggtgggtgagc 180
tcccattggt aagttttgcc ggccaagatg tcaactgatgc gttcattgct taccatcctg 240
gcactgcttg gcaatacctt gacagggttct ttactgggta ctacgttcaa gattactctg 300
tctctgagat gtccaaggac tacagaaggc tcgtctctga gttttctaag atggggttgt 360
tcaagacacc aggcaaaggg gtctactgct caatcttttt cgtgtctgtg ttgttcgctc 420
tgagtgttta cgggtgttct tactgcaaga gcacctgggc tcatctttgc tctgggttgc 480
taatgggtat gctatggctc cagagtgggt ggggtgggca tgattcttgt cactaccaag 540
ttatgcctaa ccgtaagctt aatcgctttt ttcaaatcat tgcaggaaat gtgattgctg 600
gtgttagtgt tgcattgggt aagttggacc ataacaccca tcactttgcc tghtaatagcg 660
ccaatctgga tcctgatatt cagcaccttc ctataattgc catatcccca aaatttttca 720
actcccttac atcatactat cacaactgca aaatgaccta tgatcgcgct gccagggttt 780
ttgttagctt tcagcactgg acatttttatc ctgcattggt aagcgtagg ctctatcttt 840
ttattctgtc ttttaagggt gtgttttcca acaacaaaag ggtatacaag agaagtcagg 900
aaatttttag ctatgcagct ttcttgactt ggtattctct actcctttct cgctaccca 960
attggcctga aagggtcatg tatttcacgt cctgttttag agtcgccggg ttccaacatt 1020
ggcagttcag cttgaatcac tttgcttcta atgtttacac tggtttgctt agcggtaatg 1080
attggtttca ccagcagaca aagggcacgc tcaacataac agcttctgct tgggtgggatt 1140
ggtttcatgg tggcctgcac tttcagattg agcatcatct gtttccaagg atgcctaagt 1200
gccatttcag gaaaatctca ccatttgtga acaaaacttg ccagaagcat aatttgtcct 1260
atgaaactgc taccatgtgg gaggccaata aaatggtata ctccaccctg cgtgctgtgg 1320
ctatggaagc taaggatgtt accaagccag ttcccaagaa catggtctgg gaagcaatga 1380
acactttcgg gtgaacctta tnaaacatca agtgctgtct ttcccgtaaa agcttccagt 1440
cccaatggtt cttttttttt tttttttttt t 1471

```

&lt;210&gt; 2

&lt;211&gt; 448

&lt;212&gt; PRT

&lt;213&gt; Picramnia pentandra



&lt;400&gt; 2

```

Met Glu Glu Pro Lys Lys His Ile Ser Gln Ala Asp Leu Ala Lys His
 1          5          10          15

Lys Gln Pro Gly Asp Leu Trp Ile Ser Ile Lys Gly Lys Val Tyr Asp
          20          25          30

Ile Ser Lys Trp Thr Lys Glu His Pro Gly Gly Glu Leu Pro Leu Leu
          35          40          45

Ser Phe Ala Gly Gln Asp Val Thr Asp Ala Phe Ile Ala Tyr His Pro
          50          55          60

Gly Thr Ala Trp Gln Tyr Leu Asp Arg Phe Phe Thr Gly Tyr Tyr Val
 65          70          75          80

Gln Asp Tyr Ser Val Ser Glu Met Ser Lys Asp Tyr Arg Arg Leu Val
          85          90          95

Ser Glu Phe Ser Lys Met Gly Leu Phe Lys Thr Pro Gly Lys Gly Val
          100          105          110

Tyr Cys Ser Ile Phe Phe Val Ser Val Leu Phe Ala Leu Ser Val Tyr
          115          120          125

Gly Val Leu Tyr Cys Lys Ser Thr Trp Ala His Leu Cys Ser Gly Leu
          130          135          140

Leu Met Gly Met Leu Trp Leu Gln Ser Gly Trp Val Gly His Asp Ser
          145          150          155          160

Cys His Tyr Gln Val Met Pro Asn Arg Lys Leu Asn Arg Leu Phe Gln
          165          170          175

Ile Ile Ala Gly Asn Val Ile Ala Gly Val Ser Val Ala Trp Trp Lys
          180          185          190

Leu Asp His Asn Thr His His Phe Ala Cys Asn Ser Ala Asn Leu Asp
          195          200          205

Pro Asp Ile Gln His Leu Pro Ile Ile Ala Ile Ser Pro Lys Phe Phe
          210          215          220

Asn Ser Leu Thr Ser Tyr Tyr His Asn Cys Lys Met Thr Tyr Asp Arg
          225          230          235          240

Ala Ala Arg Phe Phe Val Ser Phe Gln His Trp Thr Phe Tyr Pro Ala
          245          250          255

Leu Leu Ser Val Arg Leu Tyr Leu Phe Ile Leu Ser Phe Lys Val Val
          260          265          270

Phe Ser Asn Asn Lys Arg Val Tyr Lys Arg Ser Gln Glu Ile Leu Gly
          275          280          285

Tyr Ala Ala Phe Leu Thr Trp Tyr Ser Leu Leu Leu Ser Arg Leu Pro
          290          295          300

Asn Trp Pro Glu Arg Val Met Tyr Phe Thr Ser Cys Leu Ala Val Ala
          305          310          315          320

```

Gly Phe Gln His Trp Gln Phe Ser Leu Asn His Phe Ala Ser Asn Val  
 325 330 335  
 Tyr Thr Gly Leu Pro Ser Gly Asn Asp Trp Phe His Gln Gln Thr Lys  
 340 345 350  
 Gly Thr Leu Asn Ile Thr Ala Ser Ala Trp Trp Asp Trp Phe His Gly  
 355 360 365  
 Gly Leu His Phe Gln Ile Glu His His Leu Phe Pro Arg Met Pro Lys  
 370 375 380  
 Cys His Phe Arg Lys Ile Ser Pro Ile Val Asn Lys Leu Cys Gln Lys  
 385 390 395 400  
 His Asn Leu Ser Tyr Glu Thr Ala Thr Met Trp Glu Ala Asn Lys Met  
 405 410 415  
 Val Tyr Ser Thr Leu Arg Ala Val Ala Met Glu Ala Lys Asp Val Thr  
 420 425 430  
 Lys Pro Val Pro Lys Asn Met Val Trp Glu Ala Met Asn Thr Phe Gly  
 435 440 445

<210> 3  
 <211> 1764  
 <212> DNA  
 <213> Zea mays

<400> 3  
 gcacgagctc cctctctctc cccaatcctc cccgcctccc cctaccaaatt cagcaccacc 60  
 caaggcgcat cegagccacg gccgcgcaat gccgcctct gtcgatgcaa tgccggcccc 120  
 cggcgacgcc gcgggcgccg gcgacgtgcg catgatctcc tccaaggagc tccgcgtca 180  
 cgcttcgcgc gacgacctct ggatctccat ctccggcgac gtgtacgacg tcacgccctg 240  
 gtcccccac caccggggcg gcgacctccc gcttctcacc ctggcggggc aggacgccac 300  
 cgacgccttc ggcgcctacc accgccttc ggcgcgcccg ctctccgcgc gcttcttcgt 360  
 tggcgccctc tccgactacg ccgtctcccc cgcgtccgcg gactaccgcc gcctcctcgc 420  
 gcagctatcc tccgcgggcc tcttcgaacg cgtcggcccc accccaagg tccagctcgt 480  
 cctgatggcc gtctcttct acgcgcgct gtacctcgtc ctgcgatgcg ccagcgctg 540  
 ggcgcacctc ctgcggggg gtctcattgg ctctcgtctg atccagtcg gctggatggg 600  
 ccacgactcg ggcaccacc gcatcaccgg ccatccggtc ctgcaccgcg tcgtgcaggt 660  
 gctctccggg aactgcctca ccggcctcag catcgcttg tggagtgta accacaacac 720  
 gcaccacatc gcctgcaaca gcctggacca tgaccgggac ctccagcaca tgccgctctt 780  
 tgccgtctcc cccaagctgt tcggcaacat atggtcctac ttctaccaac ggaccctggc 840  
 gttcgatgcc gctcgaat tcttcacag ctaccagcac tggaccttct acccgtaat 900  
 gtgcacgcc aggataaatc ttctcgcgca gtccgcctg ttcgttctca cggagaagag 960  
 ggtgccgcag cgggttgctt agatcgcggg ggtcgccaca ttctgggctt ggtaccgctt 1020  
 gctggtggct tccctgccga attggtggga gaggtcgcg tttgtgcttt tcagcttcac 1080  
 catctgcggg attcagcacg tccaattctg cctgaaccac ttctcgtccg acgtgtatgt 1140  
 cgggccaccc aagggcaatg actggtttga gaagcagacg gcaggcacgc tcgacatcct 1200  
 gtgctctcct tggatggatt ggttccacgg tggcctgcag ttccagattg agcaccatct 1260  
 gtttccccgc ctacctcggg gccaccttcg caaggttgca ccggcgtcc gcgaccttg 1320  
 caagaagcat gggctcactt attctgcagc cacattctgg ggtgcaaagtg tgcttacatg 1380  
 gaagacactc agggctgctg cattgcaggc caggaccgct acaagtgggtg gtgtccgaa 1440  
 gaatttggtg tgggaggtg tgaacaccca tggataaatg ggatgaagat acgggctaag 1500  
 ggcaacttct ggtgttcagc ttggtgcccc tgtgattgtc tggatgcctt tcagttattt 1560  
 agagatatgg atcattcaac ctgcctgagt caggttgga ttttcgtgtt gacaagtggc 1620  
 tgtctatcca gttggagagt tcatgcttca atagctcgtt tggtcacggg atgttctgtt 1680

ctccctatca cggtaactat atgatgatga tccttgcttt aattcatgaa cacttgtttc 1740  
 aagattaaaa aaaaaaaaaa aaaa 1764

<210> 4  
 <211> 462  
 <212> PRT  
 <213> Zea mays

<400> 4  
 Met Pro Pro Ser Val Asp Ala Met Pro Ala Pro Gly Asp Ala Ala Gly  
 1 5 10 15  
 Ala Gly Asp Val Arg Met Ile Ser Ser Lys Glu Leu Arg Ala His Ala  
 20 25 30  
 Ser Ala Asp Asp Leu Trp Ile Ser Ile Ser Gly Asp Val Tyr Asp Val  
 35 40 45  
 Thr Pro Trp Leu Pro His His Pro Gly Gly Asp Leu Pro Leu Leu Thr  
 50 55 60  
 Leu Ala Gly Gln Asp Ala Thr Asp Ala Phe Ala Ala Tyr His Pro Pro  
 65 70 75 80  
 Ser Ala Arg Pro Leu Leu Arg Arg Phe Phe Val Gly Arg Leu Ser Asp  
 85 90 95  
 Tyr Ala Val Ser Pro Ala Ser Ala Asp Tyr Arg Arg Leu Leu Ala Gln  
 100 105 110  
 Leu Ser Ser Ala Gly Leu Phe Glu Arg Val Gly Pro Thr Pro Lys Val  
 115 120 125  
 Gln Leu Val Leu Met Ala Val Leu Phe Tyr Ala Ala Leu Tyr Leu Val  
 130 135 140  
 Leu Ala Cys Ala Ser Ala Trp Ala His Leu Leu Ala Gly Gly Leu Ile  
 145 150 155 160  
 Gly Phe Val Trp Ile Gln Ser Gly Trp Met Gly His Asp Ser Gly His  
 165 170 175  
 His Arg Ile Thr Gly His Pro Val Leu Asp Arg Val Val Gln Val Leu  
 180 185 190  
 Ser Gly Asn Cys Leu Thr Gly Leu Ser Ile Ala Trp Trp Lys Cys Asn  
 195 200 205  
 His Asn Thr His His Ile Ala Cys Asn Ser Leu Asp His Asp Pro Asp  
 210 215 220  
 Leu Gln His Met Pro Leu Phe Ala Val Ser Pro Lys Leu Phe Gly Asn  
 225 230 235 240  
 Ile Trp Ser Tyr Phe Tyr Gln Arg Thr Leu Ala Phe Asp Ala Ala Ser  
 245 250 255  
 Lys Phe Phe Ile Ser Tyr Gln His Trp Thr Phe Tyr Pro Val Met Cys  
 260 265 270

```

Ile Ala Arg Ile Asn Leu Leu Ala Gln Ser Ala Leu Phe Val Leu Thr
    275                                280                                285

Glu Lys Arg Val Pro Gln Arg Leu Leu Glu Ile Ala Gly Val Ala Thr
    290                                295                                300

Phe Trp Ala Trp Tyr Pro Leu Leu Val Ala Ser Leu Pro Asn Trp Trp
    305                                310                                315                                320

Glu Arg Val Ala Phe Val Leu Phe Ser Phe Thr Ile Cys Gly Ile Gln
    325                                330                                335

His Val Gln Phe Cys Leu Asn His Phe Ser Ser Asp Val Tyr Val Gly
    340                                345                                350

Pro Pro Lys Gly Asn Asp Trp Phe Glu Lys Gln Thr Ala Gly Thr Leu
    355                                360                                365

Asp Ile Leu Cys Ser Pro Trp Met Asp Trp Phe His Gly Gly Leu Gln
    370                                375                                380

Phe Gln Ile Glu His His Leu Phe Pro Arg Leu Pro Arg Cys His Leu
    385                                390                                395                                400

Arg Lys Val Ala Pro Ala Val Arg Asp Leu Cys Lys Lys His Gly Leu
    405                                410                                415

Thr Tyr Ser Ala Ala Thr Phe Trp Gly Ala Asn Val Leu Thr Trp Lys
    420                                425                                430

Thr Leu Arg Ala Ala Ala Leu Gln Ala Arg Thr Ala Thr Ser Gly Gly
    435                                440                                445

Ala Pro Lys Asn Leu Val Trp Glu Ala Val Asn Thr His Gly
    450                                455                                460

```

```

<210> 5
<211> 880
<212> DNA
<213> Glycine max

```

```

<220>
<221> unsure
<222> (496)

```

```

<220>
<221> unsure
<222> (512)

```

```

<220>
<221> unsure
<222> (523)

```

```

<220>
<221> unsure
<222> (532)

```

```

<220>
<221> unsure
<222> (630)

```

<220>  
 <221> unsure  
 <222> (700)

<220>  
 <221> unsure  
 <222> (730)

<220>  
 <221> unsure  
 <222> (738)

<220>  
 <221> unsure  
 <222> (761)

<220>  
 <221> unsure  
 <222> (764)

<220>  
 <221> unsure  
 <222> (814)

<220>  
 <221> unsure  
 <222> (822)

<220>  
 <221> unsure  
 <222> (824)

<220>  
 <221> unsure  
 <222> (838)

<220>  
 <221> unsure  
 <222> (842)

<220>  
 <221> unsure  
 <222> (876)

<400> 5  
 ccgcctccct cctcctcccc gccttctcca cctcccaccg tctttccgac cacaccgtct 60  
 ccgcgccttc ctccgactac cgcaagctct tctccgacct ctccgcgctc aacctcttca 120  
 accgcaaggg ccacacaacc tccatcctcc tctcccttat tctcaccctt tttcctctct 180  
 ctgtctgcgg cgtcctcttc tccgacagca ctttcgtgca cgtgctttcc gctgcattga 240  
 taggctttct ctggattcag agcggtgga taggccacga ctccggccat tacaacgtga 300  
 tgctcagccg ccgcctcaac cgcgcaattc agattctctc cggcaacatt ctgcgcggaa 360  
 tcagcatcgg ctggtggaag tggaaccaca acgcccacca cattgcatgc aacagcctcg 420  
 actatgacct tgatctgcag cacatgcggg tctttgcagt ttcgtcgcgg ttcttcaatt 480  
 ccataacctc tcatttctat gggaggaagt tngagtgtga ttncattgct angttcttga 540  
 tctgctacca gcactttact ttttaccggg taatgtgtgt tgccagggtc aacttgtatc 600  
 tgcagacaat tctgctattg ttttcgaggn gaaaagtgca ggatagagct tgaacataat 660  
 ggggatcctt gtgttttgga cttggttccct ctttttagtgn cttgcctgcc aaattgggce 720  
 tgataggggn atgtttgngc ttgctagctt tgctgtttgt nccnatccag cacattcagt 780

tctggttgaa tcaccttgct gaaaatttat atgncgggca cnantgggaa tgactggntg 840  
 anaatcagac aaggggtcat tggatatctc ttgtgncct 880

<210> 6  
 <211> 253  
 <212> PRT  
 <213> Glycine max

<220>  
 <221> UNSURE  
 <222> (161)

<220>  
 <221> UNSURE  
 <222> (166)

<220>  
 <221> UNSURE  
 <222> (170)

<220>  
 <221> UNSURE  
 <222> (173)

<220>  
 <221> UNSURE  
 <222> (206)

<220>  
 <221> UNSURE  
 <222> (252)

<400> 6  
 Leu Pro Ala Phe Ser Thr Ser His Arg Leu Ser Asp His Thr Val Ser  
 1 5 10 15  
 Ala Ala Ser Ser Asp Tyr Arg Lys Leu Phe Ser Asp Leu Ser Ala Leu  
 20 25 30  
 Asn Leu Phe Asn Arg Lys Gly His Thr Thr Ser Ile Leu Leu Ser Leu  
 35 40 45  
 Ile Leu Thr Leu Phe Pro Leu Ser Val Cys Gly Val Leu Phe Ser Asp  
 50 55 60  
 Ser Thr Phe Val His Val Leu Ser Ala Ala Leu Ile Gly Phe Leu Trp  
 65 70 75 80  
 Ile Gln Ser Gly Trp Ile Gly His Asp Ser Gly His Tyr Asn Val Met  
 85 90 95  
 Leu Ser Arg Arg Leu Asn Arg Ala Ile Gln Ile Leu Ser Gly Asn Ile  
 100 105 110  
 Leu Ala Gly Ile Ser Ile Gly Trp Trp Lys Trp Asn His Asn Ala His  
 115 120 125  
 His Ile Ala Cys Asn Ser Leu Asp Tyr Asp Pro Asp Leu Gln His Met  
 130 135 140

Pro Val Phe Ala Val Ser Ser Arg Phe Phe Asn Ser Ile Thr Ser His  
 145 150 155 160

Xaa Tyr Gly Arg Lys Xaa Glu Phe Asp Xaa Ile Ala Xaa Phe Leu Ile  
 165 170 175

Cys Tyr Gln His Phe Thr Phe Tyr Pro Val Met Cys Val Ala Arg Val  
 180 185 190

Asn Leu Tyr Leu Gln Thr Ile Leu Leu Leu Phe Ser Arg Xaa Lys Val  
 195 200 205

Gln Asp Arg Ala Leu Asn Ile Met Gly Ile Leu Val Phe Trp Thr Trp  
 210 215 220

Phe Leu Phe Leu Leu Ala Leu Leu Phe Val Pro Ile Gln His Ile Gln  
 225 230 235 240

Phe Trp Leu Asn His Leu Ala Glu Asn Leu Tyr Xaa Gly  
 245 250

<210> 7  
 <211> 1934  
 <212> DNA  
 <213> Glycine max

<400> 7

gcacgagcac	acaagtaaaa	ccttagagag	agagagagag	agagagagag	ggtaaaaggg	60
tattagatcc	ttgaaccaga	tcaaatcatc	aaaatctctg	tctatggggg	tgtgaaaaca	120
acaatcacat	tgttgttgct	gtaaaagggg	attccttagt	ctagatttgt	ttgcattgca	180
gaagaaccag	ataccccatt	attgttctta	tctatctatc	tgtctatatt	ctatttatct	240
tcttattgta	gttctcattg	tgtctgattt	cagtgtattg	tgttgttttt	ggttaacaca	300
agcaatggag	gttgttgaga	aggagaagaa	gtacataacc	tcagaggagc	tgaaggggtca	360
caacaaggag	ggagatttat	ggatctcaat	tcaaggtaag	gtgtacaatg	tctcagattg	420
ggccaaggag	caccctgggt	gtgatgttcc	aatctcaaac	cttgctggcc	aggatgtcac	480
tgatgcattc	atagcatacc	atcctggcac	agcatgggtc	caccttgaaa	aattcttcac	540
tggtaccac	ctcagtgact	tcaaggtctc	tgaggtgtcc	aaagactaca	gaaagcttgc	600
atctgagttc	tcaaaattgg	gtctttttga	caccaaaggg	catgtcactt	catgcaccct	660
tgcactctgt	gctgttatgt	tctcatttgt	actctatggg	gttctgaggt	gcactagtgt	720
gtgggctcat	ttgggttcag	gcatgctctt	agggttgctt	tggatgcaaa	gtgcttatgt	780
gggccatgat	tctggccact	atgtggttat	gacaaccaat	ggtttcaaca	aggttgcaca	840
gatcctctct	gggaactgct	tgaccgggat	aagcattgct	tgggtggaag	ggactcacia	900
tgctcaccac	attgcgtgca	acagccttga	ccatgaccct	gatctgcagc	acatgccggt	960
ctttgcagtt	tcgtcgcggt	tcttcaattc	cataacctct	catttctatg	ggaggaagtt	1020
ggagtttgat	ttcattgcta	ggttcttgat	ctgctaccag	cactttactt	tttaccgggt	1080
aatgtgtggt	gccagggtca	acttgtatct	gcagacaatt	ctgctattgt	tttcgaggcg	1140
aaaagtgcag	gatagagcct	tgaacataat	gggatccctt	gtgttttgga	cttggttccc	1200
tcttttagtg	tcttgccctgc	caaattggcc	tgagagggtt	atgtttgtgc	ttgctagctt	1260
tgctgtttgt	tccatccagc	acattcagtt	ctgtttgaat	cactttgctg	caaattgata	1320
tgctcggcca	ccgagtggga	atgactgggt	tgagaagcag	acaagtggta	cattggatat	1380
ctcttggtgc	tcttcgatgg	attgggtttt	cgggtggctt	cagtttcagc	ttgagcatca	1440
tttggtttcca	aggctacctc	ggtgccaat	gaggaagatt	tcgcctttgg	ttagtgcact	1500
ttgcaagaag	cataattttgc	cttataggag	cttgtcattt	tgggaggcca	atcagtgga	1560
aattaggacc	ctcaggactg	ctgccctaca	agctagggac	ttaacaaacc	ctgcccctaa	1620
gaatttggtg	tgggaagctg	ttaataccca	tggctgaggc	atgttgagtt	tttagagttt	1680
aggattttgt	caaggtcttt	tttttttttg	ttttctcttt	aaaaagaaaa	aaaattctca	1740
tgtgtatttt	gctagccccc	acttttccag	attgggcttt	gaatttaact	ttttgttagg	1800
tgtgggtgtac	aaatggatgg	tgatccagat	gttactgcag	ttcatgtgct	ttgcatcaat	1860
acaaattcat	atcatgtatg	ccaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	1920
aaaaaaaaaa	aaaa					1934

<210> 8  
 <211> 450  
 <212> PRT  
 <213> Glycine max

<400> 8  
 Met Glu Val Val Glu Lys Glu Lys Lys Tyr Ile Thr Ser Glu Glu Leu  
 1 5 10 15  
 Lys Gly His Asn Lys Glu Gly Asp Leu Trp Ile Ser Ile Gln Gly Lys  
 20 25 30  
 Val Tyr Asn Val Ser Asp Trp Val Lys Glu His Pro Gly Gly Asp Val  
 35 40 45  
 Pro Ile Ser Asn Leu Ala Gly Gln Asp Val Thr Asp Ala Phe Ile Ala  
 50 55 60  
 Tyr His Pro Gly Thr Ala Trp Ser His Leu Glu Lys Phe Phe Thr Gly  
 65 70 75 80  
 Tyr His Leu Ser Asp Phe Lys Val Ser Glu Val Ser Lys Asp Tyr Arg  
 85 90 95  
 Lys Leu Ala Ser Glu Phe Ser Lys Leu Gly Leu Phe Asp Thr Lys Gly  
 100 105 110  
 His Val Thr Ser Cys Thr Leu Ala Ser Val Ala Val Met Phe Leu Ile  
 115 120 125  
 Val Leu Tyr Gly Val Leu Arg Cys Thr Ser Val Trp Ala His Leu Gly  
 130 135 140  
 Ser Gly Met Leu Leu Gly Leu Leu Trp Met Gln Ser Ala Tyr Val Gly  
 145 150 155 160  
 His Asp Ser Gly His Tyr Val Val Met Thr Thr Asn Gly Phe Asn Lys  
 165 170 175  
 Val Ala Gln Ile Leu Ser Gly Asn Cys Leu Thr Gly Ile Ser Ile Ala  
 180 185 190  
 Trp Trp Lys Trp Thr His Asn Ala His His Ile Ala Cys Asn Ser Leu  
 195 200 205  
 Asp His Asp Pro Asp Leu Gln His Met Pro Val Phe Ala Val Ser Ser  
 210 215 220  
 Arg Phe Phe Asn Ser Ile Thr Ser His Phe Tyr Gly Arg Lys Leu Glu  
 225 230 235 240  
 Phe Asp Phe Ile Ala Arg Phe Leu Ile Cys Tyr Gln His Phe Thr Phe  
 245 250 255  
 Tyr Pro Val Met Cys Val Ala Arg Val Asn Leu Tyr Leu Gln Thr Ile  
 260 265 270  
 Leu Leu Leu Phe Ser Arg Arg Lys Val Gln Asp Arg Ala Leu Asn Ile  
 275 280 285



Met Gly Ile Leu Val Phe Trp Thr Trp Phe Pro Leu Leu Val Ser Cys  
 290 295 300

Leu Pro Asn Trp Pro Glu Arg Val Met Phe Val Leu Ala Ser Phe Ala  
 305 310 315 320

Val Cys Ser Ile Gln His Ile Gln Phe Cys Leu Asn His Phe Ala Ala  
 325 330 335

Asn Val Tyr Val Gly Pro Pro Ser Gly Asn Asp Trp Phe Glu Lys Gln  
 340 345 350

Thr Ser Gly Thr Leu Asp Ile Ser Cys Ala Ser Ser Met Asp Trp Phe  
 355 360 365

Phe Gly Gly Leu Gln Phe Gln Leu Glu His His Leu Phe Pro Arg Leu  
 370 375 380

Pro Arg Cys Gln Leu Arg Lys Ile Ser Pro Leu Val Ser Asp Leu Cys  
 385 390 395 400

Lys Lys His Asn Leu Pro Tyr Arg Ser Leu Ser Phe Trp Glu Ala Asn  
 405 410 415

Gln Trp Thr Ile Arg Thr Leu Arg Thr Ala Ala Leu Gln Ala Arg Asp  
 420 425 430

Leu Thr Asn Pro Ala Pro Lys Asn Leu Leu Trp Glu Ala Val Asn Thr  
 435 440 445

His Gly  
 450

<210> 9  
 <211> 1972  
 <212> DNA  
 <213> Triticum aestivum

<400> 9

gcacgagctc	cctaacaaac	ctcogttgct	gttttaagat	ccgatctccc	cttccccct	60
cccctccctt	cctcctgagt	cctgaccacc	cctcctcgcg	ctccagctaa	atccacgcca	120
ccgatggccc	gcacgggctt	cgcggaagca	acggcgccgg	aagccgacgc	aatgccggcc	180
gccagcaagg	acgccgccga	cgtecgcatg	atctccacca	aggagctgca	ggcgacgct	240
gccgcggacg	acctctggat	ctccatctcc	ggggacgtct	acgacgtcac	gcggtggctg	300
cgccaccacc	cgggcggcga	ggtcccgcgc	atcacccctg	ccggccagga	cgccaccgac	360
gccttcatgg	cctaccaccc	gcccctcggt	cgcccgtctc	tccgcccgtt	cttcgtcggc	420
cgccctctcc	actacaccgt	cccccccgcc	tccgcccact	tccgcccgtt	cctcgcgag	480
ctctcctccg	cgggcctctt	cgagcgcgct	ggccacaccc	ccaagttcct	gctcgctgca	540
atgtcggtgc	tcttctgcat	cgccctctac	tgcgtcctcg	cctgctccag	caccggggcc	600
cacatgttcg	ccggcggcct	cattggcttc	atctggatcc	agtcgggctg	gattggccat	660
gactccggcc	accaccaa	caccaggcac	cccgcgtca	accgctcct	gcagggtggtc	720
tccgggaact	gcctcaccgg	cctcggcac	gcttggtgga	agttcaacca	caacacacac	780
cacatctcct	gcaacagcct	cgaccatgac	ccggacctcc	agcacttgcc	gctcttcgcg	840
gtttccacca	agctcttcaa	caacctttgg	tgggtctgct	acgagcgcac	cttggtgttt	900
gatgccatat	ccaagttctt	cgtcagctac	cagcactgga	cattctaccc	ggtgatggga	960
tttgcaagga	taaattctct	tgtgcagtca	atcgtgttcc	tgatcacgca	aaagaaggtg	1020
cggcagcggt	ggctggagat	cgccggaggt	gcagcggtct	gggtttggta	ccccttgctg	1080
gtctcttgcc	tgccgaattg	gtgggagagg	gttgcttttg	tgcttgcaag	ctttgtgatc	1140
acgggggattc	agcatgttca	gttctgcctg	aaccacttct	catccgctgt	gtatgttggg	1200

```

ccaccaaagg ggaacgactg gtttgagagg caaacagcgg gcacacttga tatcaagtgc 1260
tccccgtgga tggattggtt ccatggtggt ctgcagttcc aggttgaaaca ccatttgttt 1320
cctcgctgc ctcgctgcca ctataggatg gtcgcgccga ttgtgcgtga cctttgcaag 1380
aagcatgggc tgtcttatgg tgccgccaca ttctgggagg caaatgtaat gacatggaag 1440
acgctaaggg ctgcagcatt gcaggccagg gaagccacta ctggagctgc tccaaagaat 1500
ctggtctggg aagctttgaa cactcatgga tgactgggat caggactgga gtatgagaca 1560
attgtaagcg tcgagccttg cgtgcatgca gttatctgat tgcttctcga ttgcgtagag 1620
atattgatcc ttttagctgt tggaatcgtg ttggattttt cgtgttgcca ggtgactatc 1680
tttgacgttc aatcgtgggt tcatgcttca gttgtgtact tgtacaccat atttagattg 1740
ttgggttctc cctatcatgg taactacatc aatagtactt gatttacatc ataaaatccg 1800
tggcttatct ttacatccat ttcattttgc ttgcaagttc atgaaactgt aaactcaatt 1860
gatggtttgt agcgtgtata tcctgctgct atggcagctt gaactgcatt ttgggaacat 1920
gacgattcca ataataaacg ttttagacatt ttctaaaaaa aaaaaaaaaa aa 1972

```

```

<210> 10
<211> 469
<212> PRT
<213> Triticum aestivum

```

```

<400> 10
Met Ala Arg Thr Gly Leu Ala Asp Ala Thr Ala Pro Glu Ala Asp Ala
  1                      5                      10                      15

Met Pro Ala Ala Ser Lys Asp Ala Ala Asp Val Arg Met Ile Ser Thr
          20                      25                      30

Lys Glu Leu Gln Ala His Ala Ala Ala Asp Asp Leu Trp Ile Ser Ile
          35                      40                      45

Ser Gly Asp Val Tyr Asp Val Thr Pro Trp Leu Arg His His Pro Gly
  50                      55                      60

Gly Glu Val Pro Leu Ile Thr Leu Ala Gly Gln Asp Ala Thr Asp Ala
  65                      70                      75                      80

Phe Met Ala Tyr His Pro Pro Ser Val Arg Pro Leu Leu Arg Arg Phe
          85                      90                      95

Phe Val Gly Arg Leu Ser Asp Tyr Thr Val Pro Pro Ala Ser Ala Asp
  100                      105                      110

Phe Arg Arg Leu Leu Ala Gln Leu Ser Ser Ala Gly Leu Phe Glu Arg
  115                      120                      125

Val Gly His Thr Pro Lys Phe Leu Leu Val Ala Met Ser Val Leu Phe
  130                      135                      140

Cys Ile Ala Leu Tyr Cys Val Leu Ala Cys Ser Ser Thr Gly Ala His
  145                      150                      155                      160

Met Phe Ala Gly Gly Leu Ile Gly Phe Ile Trp Ile Gln Ser Gly Trp
          165                      170                      175

Ile Gly His Asp Ser Gly His His Gln Ile Thr Arg His Pro Ala Leu
  180                      185                      190

Asn Arg Leu Leu Gln Val Val Ser Gly Asn Cys Leu Thr Gly Leu Gly
  195                      200                      205

```

Ile Ala Trp Trp Lys Phe Asn His Asn Thr His His Ile Ser Cys Asn  
 210 215 220  
 Ser Leu Asp His Asp Pro Asp Leu Gln His Leu Pro Leu Phe Ala Val  
 225 230 235 240  
 Ser Thr Lys Leu Phe Asn Asn Leu Trp Ser Val Cys Tyr Glu Arg Thr  
 245 250 255  
 Leu Ala Phe Asp Ala Ile Ser Lys Phe Phe Val Ser Tyr Gln His Trp  
 260 265 270  
 Thr Phe Tyr Pro Val Met Gly Phe Ala Arg Ile Asn Leu Leu Val Gln  
 275 280 285  
 Ser Ile Val Phe Leu Ile Thr Gln Lys Lys Val Arg Gln Arg Trp Leu  
 290 295 300  
 Glu Ile Ala Gly Val Ala Ala Phe Trp Val Trp Tyr Pro Leu Leu Val  
 305 310 315 320  
 Ser Cys Leu Pro Asn Trp Trp Glu Arg Val Ala Phe Val Leu Ala Ser  
 325 330 335  
 Phe Val Ile Thr Gly Ile Gln His Val Gln Phe Cys Leu Asn His Phe  
 340 345 350  
 Ser Ser Ala Val Tyr Val Gly Pro Pro Lys Gly Asn Asp Trp Phe Glu  
 355 360 365  
 Arg Gln Thr Ala Gly Thr Leu Asp Ile Lys Cys Ser Pro Trp Met Asp  
 370 375 380  
 Trp Phe His Gly Gly Leu Gln Phe Gln Val Glu His His Leu Phe Pro  
 385 390 395 400  
 Arg Leu Pro Arg Cys His Tyr Arg Met Val Ala Pro Ile Val Arg Asp  
 405 410 415  
 Leu Cys Lys Lys His Gly Leu Ser Tyr Gly Ala Ala Thr Phe Trp Glu  
 420 425 430  
 Ala Asn Val Met Thr Trp Lys Thr Leu Arg Ala Ala Ala Leu Gln Ala  
 435 440 445  
 Arg Glu Ala Thr Thr Gly Ala Ala Pro Lys Asn Leu Val Trp Glu Ala  
 450 455 460  
 Leu Asn Thr His Gly  
 465

<210> 11  
 <211> 448  
 <212> PRT  
 <213> Borago officinalis

<400> 11  
 Met Ala Ala Gln Ile Lys Lys Tyr Ile Thr Ser Asp Glu Leu Lys Asn  
 1 5 10 15

His Asp Lys Pro Gly Asp Leu Trp Ile Ser Ile Gln Gly Lys Ala Tyr  
 20 25 30  
 Asp Val Ser Asp Trp Val Lys Asp His Pro Gly Gly Ser Phe Pro Leu  
 35 40 45  
 Lys Ser Leu Ala Gly Gln Glu Val Thr Asp Ala Phe Val Ala Phe His  
 50 55 60  
 Pro Ala Ser Thr Trp Lys Asn Leu Asp Lys Phe Phe Thr Gly Tyr Tyr  
 65 70 75 80  
 Leu Lys Asp Tyr Ser Val Ser Glu Val Ser Lys Asp Tyr Arg Lys Leu  
 85 90 95  
 Val Phe Glu Phe Ser Lys Met Gly Leu Tyr Asp Lys Lys Gly His Ile  
 100 105 110  
 Met Phe Ala Thr Leu Cys Phe Ile Ala Met Leu Phe Ala Met Ser Val  
 115 120 125  
 Tyr Gly Val Leu Phe Cys Glu Gly Val Leu Val His Leu Phe Ser Gly  
 130 135 140  
 Cys Leu Met Gly Phe Leu Trp Ile Gln Ser Gly Trp Ile Gly His Asp  
 145 150 155 160  
 Ala Gly His Tyr Met Val Val Ser Asp Ser Arg Leu Asn Lys Phe Met  
 165 170 175  
 Gly Ile Phe Ala Ala Asn Cys Leu Ser Gly Ile Ser Ile Gly Trp Trp  
 180 185 190  
 Lys Trp Asn His Asn Ala His His Ile Ala Cys Asn Ser Leu Glu Tyr  
 195 200 205  
 Asp Pro Asp Leu Gln Tyr Ile Pro Phe Leu Val Val Ser Ser Lys Phe  
 210 215 220  
 Phe Gly Ser Leu Thr Ser His Phe Tyr Glu Lys Arg Leu Thr Phe Asp  
 225 230 235 240  
 Ser Leu Ser Arg Phe Phe Val Ser Tyr Gln His Trp Thr Phe Tyr Pro  
 245 250 255  
 Ile Met Cys Ala Ala Arg Leu Asn Met Tyr Val Gln Ser Leu Ile Met  
 260 265 270  
 Leu Leu Thr Lys Arg Asn Val Ser Tyr Arg Ala His Glu Leu Leu Gly  
 275 280 285  
 Cys Leu Val Phe Ser Ile Trp Tyr Pro Leu Leu Val Ser Cys Leu Pro  
 290 295 300  
 Asn Trp Gly Glu Arg Ile Met Phe Val Ile Ala Ser Leu Ser Val Thr  
 305 310 315 320  
 Gly Met Gln Gln Val Gln Phe Ser Leu Asn His Phe Ser Ser Ser Val  
 325 330 335

Tyr Val Gly Lys Pro Lys Gly Asn Asn Trp Phe Glu Lys Gln Thr Asp  
 340 345 350  
 Gly Thr Leu Asp Ile Ser Cys Pro Pro Trp Met Asp Trp Phe His Gly  
 355 360 365  
 Gly Leu Gln Phe Gln Ile Glu His His Leu Phe Pro Lys Met Pro Arg  
 370 375 380  
 Cys Asn Leu Arg Lys Ile Ser Pro Tyr Val Ile Glu Leu Cys Lys Lys  
 385 390 395 400  
 His Asn Leu Pro Tyr Asn Tyr Ala Ser Phe Ser Lys Ala Asn Glu Met  
 405 410 415  
 Thr Leu Arg Thr Leu Arg Asn Thr Ala Leu Gln Ala Arg Asp Ile Thr  
 420 425 430  
 Lys Pro Leu Pro Lys Asn Leu Val Trp Glu Ala Leu His Thr His Gly  
 435 440 445  
 <210> 12  
 <211> 469  
 <212> PRT  
 <213> Triticum aestivum  
 <400> 12  
 Met Ala Arg Thr Gly Leu Ala Asp Ala Thr Ala Pro Glu Ala Asp Ala  
 1 5 10 15  
 Met Pro Ala Ala Ser Lys Asp Ala Ala Asp Val Arg Met Ile Ser Thr  
 20 25 30  
 Lys Glu Leu Gln Ala His Ala Ala Ala Asp Asp Leu Trp Ile Ser Ile  
 35 40 45  
 Ser Gly Asp Val Tyr Asp Val Thr Pro Trp Leu Arg His His Pro Gly  
 50 55 60  
 Gly Glu Val Pro Leu Ile Thr Leu Ala Gly Gln Asp Ala Thr Asp Ala  
 65 70 75 80  
 Phe Met Ala Tyr His Pro Pro Ser Val Arg Pro Leu Leu Arg Arg Phe  
 85 90 95  
 Phe Val Gly Arg Leu Thr Asp Tyr Thr Val Pro Pro Ala Ser Ala Asp  
 100 105 110  
 Phe Arg Arg Leu Leu Ala Gln Leu Ser Ser Ala Gly Leu Phe Glu Arg  
 115 120 125  
 Val Gly His Thr Pro Lys Phe Leu Leu Val Ala Met Ser Val Leu Phe  
 130 135 140  
 Cys Ile Ala Leu Tyr Cys Val Leu Ala Cys Ser Ser Thr Gly Ala His  
 145 150 155 160  
 Met Phe Ala Gly Gly Leu Ile Gly Phe Ile Trp Ile Gln Ser Gly Trp  
 165 170 175

Ile Gly His Asp Ser Gly His His Gln Ile Thr Arg His Pro Ala Leu  
 180 185 190  
 Asn Arg Leu Leu Gln Val Val Ser Gly Asn Cys Leu Thr Gly Leu Gly  
 195 200 205  
 Ile Ala Trp Trp Lys Phe Asn His Asn Thr His His Ile Ser Cys Asn  
 210 215 220  
 Ser Leu Asp His Asp Pro Asp Leu Gln His Leu Pro Leu Phe Ala Val  
 225 230 235 240  
 Ser Thr Lys Leu Phe Asn Asn Leu Trp Ser Val Cys Tyr Glu Arg Thr  
 245 250 255  
 Leu Ala Phe Asp Ala Ile Ser Lys Phe Phe Val Ser Tyr Gln His Trp  
 260 265 270  
 Thr Phe Tyr Pro Val Met Gly Phe Ala Arg Ile Asn Leu Leu Val Gln  
 275 280 285  
 Ser Ile Val Phe Leu Ile Thr Gln Lys Lys Val Arg Gln Arg Trp Leu  
 290 295 300  
 Glu Ile Ala Gly Val Ala Ala Phe Trp Val Trp Tyr Pro Leu Leu Val  
 305 310 315 320  
 Ser Cys Leu Pro Asn Trp Trp Glu Arg Val Ala Phe Val Leu Ala Ser  
 325 330 335  
 Phe Val Ile Thr Gly Ile Gln His Val Gln Phe Cys Leu Asn His Phe  
 340 345 350  
 Ser Ser Ala Val Tyr Val Gly Pro Pro Lys Gly Asn Asp Trp Phe Glu  
 355 360 365  
 Arg Gln Thr Ala Gly Thr Leu Asp Ile Lys Cys Ser Pro Trp Met Asp  
 370 375 380  
 Trp Phe His Gly Gly Leu Gln Phe Gln Val Glu His His Leu Phe Pro  
 385 390 395 400  
 Arg Leu Pro Arg Cys His Tyr Arg Met Val Ala Pro Ile Val Arg Asp  
 405 410 415  
 Leu Cys Lys Lys His Gly Leu Ser Tyr Gly Ala Ala Thr Phe Trp Glu  
 420 425 430  
 Ala Asn Val Met Thr Trp Lys Thr Leu Arg Ala Ala Ala Leu Gln Ala  
 435 440 445  
 Arg Glu Ala Thr Thr Gly Ala Ala Pro Lys Asn Leu Val Trp Glu Ala  
 450 455 460  
 Leu Asn Thr His Gly  
 465

&lt;210&gt; 13

&lt;211&gt; 458

&lt;212&gt; PRT

&lt;213&gt; Helianthus annuus

&lt;400&gt; 13

```

Met Val Ser Pro Ser Ile Glu Val Leu Asn Ser Ile Ala Asp Gly Lys
 1           5           10           15

Lys Tyr Ile Thr Ser Lys Glu Leu Lys Lys His Asn Asn Pro Asn Asp
          20           25           30

Leu Trp Ile Ser Ile Leu Gly Lys Val Tyr Asn Val Thr Glu Trp Ala
          35           40           45

Lys Glu His Pro Gly Gly Asp Ala Pro Leu Ile Asn Leu Ala Gly Gln
          50           55           60

Asp Val Thr Asp Ala Phe Ile Ala Phe His Pro Gly Thr Ala Trp Lys
          65           70           75           80

His Leu Asp Lys Leu Phe Thr Gly Tyr His Leu Lys Asp Tyr Gln Val
          85           90           95

Ser Asp Ile Ser Arg Asp Tyr Arg Lys Leu Ala Ser Glu Phe Ala Lys
          100          105          110

Ala Gly Met Phe Glu Lys Lys Gly His Gly Val Ile Tyr Ser Leu Cys
          115          120          125

Phe Val Ser Leu Leu Leu Ser Ala Cys Val Tyr Gly Val Leu Tyr Ser
          130          135          140

Gly Ser Phe Trp Ile His Met Leu Ser Gly Ala Ile Leu Gly Leu Ala
          145          150          155          160

Trp Met Gln Ile Ala Tyr Leu Gly His Asp Ala Gly His Tyr Gln Met
          165          170          175

Met Ala Thr Arg Gly Trp Asn Lys Phe Ala Gly Ile Phe Ile Gly Asn
          180          185          190

Cys Ile Thr Gly Ile Ser Ile Ala Trp Trp Lys Trp Thr His Asn Ala
          195          200          205

His His Ile Ala Cys Asn Ser Leu Asp Tyr Asp Pro Asp Leu Gln His
          210          215          220

Leu Pro Met Leu Ala Val Ser Ser Lys Leu Phe Asn Ser Ile Thr Ser
          225          230          235          240

Val Phe Tyr Gly Arg Gln Leu Thr Phe Asp Pro Leu Ala Arg Phe Phe
          245          250          255

Val Ser Tyr Gln His Tyr Leu Tyr Tyr Pro Ile Met Cys Val Ala Arg
          260          265          270

Val Asn Leu Tyr Leu Gln Thr Ile Leu Leu Leu Ile Ser Lys Arg Lys
          275          280          285

Ile Pro Asp Arg Gly Leu Asn Ile Leu Gly Thr Leu Ile Phe Trp Thr
          290          295          300

```

Trp Phe Pro Leu Leu Val Ser Arg Leu Pro Asn Trp Pro Glu Arg Val  
 305 310 315 320  
 Ala Phe Val Leu Val Ser Phe Cys Val Thr Gly Ile Gln His Ile Gln  
 325 330 335  
 Phe Thr Leu Asn His Phe Ser Gly Asp Val Tyr Val Gly Pro Pro Lys  
 340 345 350  
 Gly Asp Asn Trp Phe Glu Lys Gln Thr Arg Gly Thr Ile Asp Ile Ala  
 355 360 365  
 Cys Ser Ser Trp Met Asp Trp Phe Phe Gly Gly Leu Gln Phe Gln Leu  
 370 375 380  
 Glu His His Leu Phe Pro Arg Leu Pro Arg Cys His Leu Arg Ser Ile  
 385 390 395 400  
 Ser Pro Ile Cys Arg Glu Leu Cys Lys Lys Tyr Asn Leu Pro Tyr Val  
 405 410 415  
 Ser Leu Ser Phe Tyr Asp Ala Asn Val Thr Thr Leu Lys Thr Leu Arg  
 420 425 430  
 Thr Ala Ala Leu Gln Ala Arg Asp Leu Thr Asn Pro Ala Pro Gln Asn  
 435 440 445  
 Leu Ala Trp Glu Ala Phe Asn Thr His Gly  
 450 455

<210> 14

<211> 38

<212> DNA

<213> Artificial Sequence

<220>

<223> Definition of Artificial Sequence: PCR primer for 5' of pk0011.d5

<400> 14

tttgcggccg caaatcaatg gaagaagcaa agaag

35

<210> 15

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Definition of Sequence: antisense PCR primer for 3' of pk0011.d5

<400> 15

tttgcggccg ccaggattca cccgaaagtg ttc

33

<210> 16

<211> 823

<212> DNA

<213> Triticum aestivum



&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (48)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (538)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (686)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (704)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (717)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (727)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (729)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (737)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (741)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (750) .. (751)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (769)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (777)

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; (807)

&lt;400&gt; 16

```

ctccctaaca aacctccggt gctgttttaa gatccgatct ccccttcncc cctccccctcc 60
cttcctcctg agtcctgacc acccctcctc gcgctccagc taaatccacg ccaccgatgg 120
cccgcacggg cttcgcgga cgaacggcgc cgggaagccga cgcaatgccg gccgccagca 180
aggacgccgc cgacgtccgc atgatctcca ccaaggagct gcaggcgcac gccgccgcgg 240
acgacctctg gatctccatc tccggggacg tctacgacgt cacgccctgg ctgcgccacc 300
acccggggcg cgaggteccg ctcatacccc tcgccggcca ggacgccacc gacgccttca 360
tggcctacca cccgccctcc gtgcgccgcg tcctccgccg cttcttcgtc ggccgcctca 420

```

```

ccgactacac tgtccccccc gcttcgcgcg acttcgcgcg cctcctcgcg cagctctcct 480
ccgcggggcct cttcgagcgc gtcggcacac ccccaagttc ctgctcgtcg caaagtcngt 540
gctctttctgc atcggcctct actgctcctc gctgctcaa caccggggcc acatgttcgc 600
cgggggctca ttggcttata tggtcagtcg ggctggattg gcatactccg gcacacaaat 660
cacaggcacc tgcctcaacg ctctgnagtg gctcgggaat gctnacggct cggatcnctg 720
gggagtnanc acacaanaca nattctgaan ngtcacatac ctgactcana ttccgtntcg 780
ggtcacaagt ctaaaacttg catcgtnaag acttggttag cat 823

```

<210> 17  
 <211> 114  
 <212> PRT  
 <213> Triticum aestivum

<400> 17  
 Met Pro Ala Ala Ser Lys Asp Ala Ala Asp Val Arg Met Ile Ser Thr  
 1 5 10 15  
 Lys Glu Leu Gln Ala His Ala Ala Ala Asp Asp Leu Trp Ile Ser Ile  
 20 25 30  
 Ser Gly Asp Val Tyr Asp Val Thr Pro Trp Leu Arg His His Pro Gly  
 35 40 45  
 Gly Glu Val Pro Leu Ile Thr Leu Ala Gly Gln Asp Ala Thr Asp Ala  
 50 55 60  
 Phe Met Ala Tyr His Pro Pro Ser Val Arg Pro Leu Leu Arg Arg Phe  
 65 70 75 80  
 Phe Val Gly Arg Leu Thr Asp Tyr Thr Val Pro Pro Ala Ser Ala Asp  
 85 90 95  
 Phe Arg Arg Leu Leu Ala Gln Leu Ser Ser Ala Gly Leu Phe Glu Arg  
 100 105 110  
 Val Gly